

**CORRECTED
VERSION***

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A61K 31/66, 39/12, 39/42, C12N 15/33, G01N 33/53, 33/533, 33/534, 33/535, 33/569</p>	<p>A1</p>	<p>(11) International Publication Number: WO 96/06619</p>
<p>(21) International Application Number: PCT/US95/10904 (22) International Filing Date: 1 September 1995 (01.09.95)</p>		<p>(43) International Publication Date: 7 March 1996 (07.03.96)</p>
<p>(30) Priority Data: 301,435 1 September 1994 (01.09.94) US</p> <p>(71)(72) Applicants and Inventors: PAUL, Prem, S. [US/US]; 4206 Arizona Circle, Ames, IA 50014 (US). MENG, Xiang- Jin [CN/US]; 725 Pammel Court, Ames, IA 50014 (US). HALBUR, Patrick [US/US]; 3211 Kingman Road, Ames, IA 50014 (US). MOROZOV, Igor [RU/US]; Apartment 156, 407 South 5th Street, Ames, IA 50010 (US). LUM, Melissa, A. [US/US]; Solvay Animal Health, Inc., 1201 Northland Drive, Mendota Heights, MN 55120-1149 (US).</p> <p>(74) Agents: LAVALLEYE, Jean-Paul et al.; Oblon, Spivak, McClelland, Maier & Neustadt, P.C., 4th floor, Crystal Square Five, 1755 Jefferson Davis Highway, Arlington, VA 22202 (US).</p>		<p>(81) Designated States: CA, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published With international search report.</p>
<p>(54) Title: POLYNUCLEIC ACIDS AND PROTEINS FROM A PORCINE REPRODUCTIVE AND RESPIRATORY SYNDROME VIRUS AND USES THEREOF</p>		
<p>(57) Abstract</p> <p>The present invention provides a purified preparation containing a polynucleic acid encoding at least one polypeptide selected from the group consisting of proteins encoded by one or more open reading frames (ORF's) of an Iowa strain of porcine reproductive and respiratory syndrome virus (PRRSV), proteins homologous with those encoded by one or more of the ORF's, antigenic regions of such proteins which are at least 5 amino acids in length and which effectively stimulate immunological protection in a porcine host against a subsequent challenge with a PRRSV isolate, and combinations thereof, in which amino acids non-essential for antigenicity may be conservatively substituted. The present invention also concerns a polypeptide encoded by such a polynucleic acid, a vaccine comprising an effective amount of such a polynucleic acid or protein, antibodies which specifically bind to such a polynucleic acid or protein; methods of producing the same; and methods of raising an effective immunological response against PRRSV, treating a pig infected by PRRSV, and detecting PRRSV.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

-1-
Description

POLYNUCLEIC ACIDS AND PROTEINS FROM A PORCINE REPRODUCTIVE AND RESPIRATORY SYNDROME VIRUS AND USES THEREOF

This is a continuation-in-part of application Serial No. 08/131,625, filed on October 5, 1993, pending, which is a continuation-in-part of application Serial No. 07/969,071, filed on October 30, 1992, now abandoned. The entire contents of application Serial No. 08/131,625, filed on October 5, 1993, are incorporated herein by reference.

Field of the Invention

The present invention concerns DNA isolated from a porcine reproductive and respiratory virus (PRRSV), a protein and/or a polypeptide encoded by the DNA, a vaccine which protects pigs from a PRRSV based on the protein or DNA, a method of protecting a pig from a PRRSV using the vaccine, a method of producing the vaccine, a method of treating a pig infected by or exposed to a PRRSV, and a method of detecting a PRRSV.

Discussion of the Background:

In recent years, North American and European swine herds have been susceptible to infection by new strains of reproductive and respiratory viruses (see A.A.S.P., September/October 1991, pp. 7-11; *The Veterinary Record*, February 1, 1992, pp. 87-89; *Ibid.*, November 30, 1991, pp. 495-496; *Ibid.*, October 26, 1991, p. 370; *Ibid.*, October 19, 1991, pp. 367-368; *Ibid.*, August 3, 1991, pp. 102-103; *Ibid.*, July 6, 1991; *Ibid.*, June 22, 1991, p. 578; *Ibid.*, June 15, 1991, p. 574; *Ibid.*, June 8, 1991, p. 536; *Ibid.*, June 1, 1991, p. 511; *Ibid.*, March 2, 1991, p. 213). Among the first of the new strains to be identified was a virus associated with the so-called Mystery Swine Disease (MSD) or "blue-eared syndrome", now known as Swine Infertility

and Respiratory Syndrome (SIRS) or Porcine Reproductive and Respiratory Syndrome (PRRS).

An MSD consisting of reproductive failure in females and respiratory disease in nursing and weaned pigs appeared in the midwestern United States in 1987 (Hill et al., *Am. Assoc. Swine Practitioner Newsletter* 4:47 (1992); Hill et al., *Proceedings Mystery Swine Disease Committee Meeting*, Denver, Colorado 29-31 (1990); Keffaber, *Am. Assoc. Swine Practitioner Newsletter* 1:1-9 (1989); Loula, *Agri-Practice* 12:23-34 (1991)). Reproductive failure was characterized by abortions, stillborn and weak-born pigs. The respiratory disease in nursing and weaned pigs was characterized by fever, labored breathing and pneumonia. A similar disease appeared in Europe in 1990 (Paton et al., *Vet. Rec.* 128:617 (1991); Wensvoort et al., *Veterinary Quarterly* 13:121-130 (1991); Blaha, *Proc. Am. Assoc. Swine Practitioners*, pp. 313-315 (1993)), and has now been recognized worldwide.

This disease has also been called porcine epidemic abortion and respiratory syndrome (PEARS), blue abortion disease, blue ear disease (U.K.), abortus blau (Netherlands), seuchenhafter spatabort der schweine (Germany), Heko-Heko disease, and in the U.S., Wabash syndrome, mystery pig disease (MPD), and swine plague (see the references cited above and Meredith, *Review of Porcine Reproductive and Respiratory Disease Syndrome*, Pig Disease Information Centre, Department of Veterinary Medicine, Madingley Road, Cambridge CB3 OES, U.K. (1992); Wensvoort et al., *Vet. Res.* 24:117-124 (1993); Paul et al., *J. Clin. Vet. Med.* 11:19-28 (1993)). In Europe, the corresponding virus has been termed "Lelystad virus."

At an international conference in May, 1992, researchers from around the world agreed to call this disease Porcine Reproductive and Respiratory Syndrome (PRRS). The disease originally appeared to be mainly a

reproductive disease during its early phases, but has now evolved primarily into a respiratory disease.

Porcine reproductive and respiratory syndrome virus (PRRSV) is a relatively recently recognized swine pathogen associated with porcine reproductive and respiratory syndrome (PRRS). PRRSV is a significant pathogen in the swine industry. PRRSV infections are common in the U.S. swine herds. Outbreaks of PRRS in England have led to cancellation of pig shows.

The symptoms of PRRS include a reluctance to eat (anorexia), a mild fever (pyrexia), cyanosis of the extremities (notably bluish ears), stillbirths, abortion, high mortality in affected litters, weak-born piglets and premature farrowing. The majority of piglets born alive to affected sows die within 48 hours. PRRS clinical signs include mild influenza-like signs, rapid respiration ("thumping"), and a diffuse interstitial pneumonitis. PRRS virus has an incubation period of about 1-2 weeks from contact with a PRRSV-infected animal. The virus appears to be an enveloped RNA arterivirus (*The Veterinary Record*, February 1, 1992). The virus has been grown successfully in pig alveolar macrophages and CL2621 cells (Benfield et al, *J. Vet. Diagn. Invest.*, 4:127-133, 1992; Collins et al, *Swine Infertility and Respiratory Syndrome/Mystery Swine Disease. Proc., Minnesota Swine Conference for Veterinarians*, pp. 200-205, 1991), and in MARC-145 cells (Joo, *PRRS: Diagnosis, Proc., Allen D. Leman Swine Conference, Veterinary Continuing Education and Extension, University of Minnesota* (1993), 20:53-55; Kim et al, *Arch. Virol.*, 133:477-483 (1993)). A successful culturing of a virus which causes SIRS has also been reported by Wensvoort et al (*Mystery Swine Disease in the Netherlands: The Isolation of Lelystad Virus. Vet. Quart.* 13:121-130, 1991).

Initially, a number of agents were incriminated in the etiology of this disease (Wensvoort et al., *Vet. Res.* 24:117-124 (1993); Woolen et al., *J. Am. Vet. Med. Assoc.* 197:600-601 (1990)). There is now a consensus that the causative agent of PRRS is an enveloped RNA virus referred to as Porcine Reproductive and Respiratory Syndrome Virus (PRRSV), reportedly of approximately 62 nm in diameter (Benfield et al., *J. Vet. Diagn. Invest.*, 4:127-133, 1992).

Virus isolates vary in their ability to replicate in continuous cell lines. Some grow readily, while others require several passages and some grow only in swine alveolar (SAM) cultures (Bautista et al., *J. Vet. Diagn. Invest.* 5:163-165, 1993; see also the Examples hereunder [particularly Table 1]).

PRRSV is a member of an Arterivirus group which includes equine arteritis virus (EAV), lactate dehydrogenase-elevating virus (LDV) and simian hemorrhagic fever virus (SHFV) (Benfield et al., 1992, *supra*; Plagemann, *Proc. Am. Assoc. Swine Practitioners*, 4:8-15 1992; Plagemann and Moennig, *Adv. Virus Res.* 41:99-192, 1992; Conzelmann et al., *Virology*, 193:329-339, 1993; Godney et al., *Virology*, 194:585-596, 1993; Meulenberg et al., *Virology*, 192:62-72, 1993). The positive-strand RNA viruses of this Arterivirus group resemble togaviruses morphologically, but are distantly related to coronaviruses and toroviruses on the basis of genome organization and gene expression (Plagemann et al., *supra*; Spaan et al., *J. Gen. Virol.* 69, 2939-2952 (1988); Strauss et al., *Annu. Rev. Biochem.* 42, 657-683 (1988); Lai, *Annu. Rev. Microbiol.* 44, 303-333 (1990); Snijder et al., *Nucleic Acid Res.* 18, 4535-4542 (1990)). The members of this group infect macrophages and contain a nested set of 5 to 7 subgenomic mRNAs in infected cells (Plagemann et al., *supra*; Meulenberg et al., *Virology*, 192, 62-72 (1993);

Conzelmann et al., *Virology*, 193, 329-339 (1993); 15, 16, 17, 18, 19).

The viral genome of European isolates has been shown to be a plus stranded RNA of about 15.1 kb (Conzelmann et al., *supra*; Meulenberg et al., *supra*), and appears to be similar in genomic organization to LDV and EAV (Meulenberg et al., *supra*). However, no serological cross-reaction has been found among PRRSV, LDV and EAV (Goyal et al., *J. Vet. Diagn. Invest.*, 5, 656-664 (1993)).

PRRSV was initially cultivated in swine alveolar macrophage (SAM) cell cultures (Pol et al., *Veterinary Quarterly*, 13:137-143, 1991; Wensvoort et al., *Veterinary Quarterly*, 13:121-130, 1991) and then in continuous cell lines CL2621 (Benfield et al., *supra*), MA-104, and MARC-145 (Joo, *Proc. Allen D. Leman Swine Conference*, pp. 53-55, 1993). The reproductive and respiratory disease has been reproduced with cell free lung filtrates (Christianson et al., *Am. J. Vet. Res.*, 53:485-488, 1992; Collins et al., *J. Vet. Diagn. Invest.*, 4:117-126, 1992; Halbur et al., *Proc. Central Veterinary Conference*, pp. 50-59, 1993), and with cell culture-propagated PRRSV (Collins et al., *supra*, and *Proc. Allen D. Leman Swine Conference*, pp. 47-48, 1993).

Eight open reading frames (also referred to herein as "ORFs" or "genes") have been identified in a European PRRSV isolate. The genes of this European isolate are organized similarly to that in coronavirus (Meulenberg et al., *supra*). A 3'-end nested set of messenger RNA has been found in PRRSV-infected cells similar to that in coronaviruses (Conzelmann et al., *supra*; Meulenberg et al., *supra*).

The ORF 1a and 1b at the 5'-half of the European PRRSV genome are predicted to encode viral RNA polymerase. The ORF's 2-6 at the 3'-half of the genome likely encode for viral membran-associated (envelope) proteins (Meulenberg et al., *supra*). ORF6 is predicted to encode the membrane

protein (M) based on its similar characteristics with the ORF 6 of EAV, ORF 2 of LDV, and the M protein of mouse hepatitis virus and infectious bronchitis virus (Meulenberg et al., *Virology* 192, 62-72 (1993); Conzelmann et al., *Virology* 193, 329-339 (1993); Murtaugh, *Proc. Allen D. Leman Swine Conference*, Minneapolis, MN, pp. 43-45 (1993); Mardassi et al., *Abstracts of Conference of Research Workers in Animal Diseases*, Chicago, IL, pp. 43 (1993)). The product of ORF 7 is extremely basic and hydrophilic, and is predicted to be the viral nucleocapsid protein (N) (Meulenberg et al., *supra*; Conzelmann et al., *supra*; Murtaugh, *supra*; Mardassi et al., *supra* and *J. Gen. Virol.*, 75:681-685 (1994)).

Although conserved epitopes have been identified between U.S. and European PRRSV isolates using monoclonal antibodies (Nelson et al., *J. Clin. Microbiol.*, 31:3184-3189, 1993), there is extensive antigenic and genetic variation both among U.S. and European isolates of PRRSV (Wensvoort et al., *J. Vet. Diagn. Invest.*, 4:134-138, 1992). European isolates are genetically closely related, as the nucleotide sequence at the 3'-half of the genome from two European PRRSV isolates is almost identical (Conzelmann et al., *supra*; Meulenberg et al., *supra*).

Although the syndrome caused by PRRSV appears to be similar in the U.S. and Europe, several recent studies have described phenotypic, antigenic, genetic and pathogenic variations among PRRSV isolates in the U.S. and in Europe (Murtaugh, *supra*; Bautista et al., *J. Vet. Diagn. Invest.*, 5, 163-165 (1993); Bautista et al., *J. Vet. Diagn. Invest.*, 5, 612-614 (1993); Wensvoort et al., *J. Vet. Diagn. Invest.*, 4, 134-138 (1992); Stevenson et al., *J. Vet. Diagn. Invest.*, 5, 432-434 (1993)). For example, the European isolates grow preferentially in SAM cultures and replicate to a very low titer in other culture systems (Wensvoort, *Vet. Res.*, 24, 117-124 (1993); Wensvoort et

al., *J. Vet. Quart.*, 13, 121-130 (1991); Wensvoort et al., *J. Vet. Diagn. Invest.*, 4, 134-138 (1992)). On the other hand, some of the U.S. isolates have been shown to replicate well in SAM as well as in the continuous cell line CL2621 (Benfield et al., *J. Vet. Diagn. Invest.*, 4, 127-133 (1992); Collins et al., *J. Vet. Diagn. Invest.*, 4, 117-126 (1992)). Thus, phenotypic differences among U.S. isolates are observed, as not all PRRSV isolates isolated on SAM can replicate on the CL2621 cell line (Bautista et al., *J. Vet. Diagn. Invest.*, 5, 163-165 (1993)).

A high degree of regional antigenic variation among PRRSV isolates may exist. Four European isolates were found to be closely related antigenically, but these European isolates differed antigenically from U.S. isolates. Further, three U.S. isolates were shown to differ antigenically from each other (Wensvoort et al., *J. Vet. Diagn. Invest.*, 4, 134-138 (1992)). Animals seropositive for European isolates were found to be negative for U.S. isolate VR 2332 (Bautista et al., *J. Vet. Diagn. Invest.*, 5, 612-614 (1993)).

U.S. PRRSV isolates differ genetically at least in part from European isolates (Conzelmann et al., *supra*; Meulenberg et al., *supra*; Murtaugh et al., *Proc. Allen D. Lemman Conference*, pp. 43-45, 1993). The genetic differences between U.S. and European isolates are striking, especially since they are considered to be the same virus (Murtaugh, *supra*). Similar observations were also reported when comparing the Canadian isolate IAF-exp91 and another U.S. isolate VR 2332 with LV (Murtaugh, *supra*; Mardassi, *supra*). However, the 3' terminal 5 kb nucleotide sequences of two European isolates are almost identical (Conzelmann et al., *supra*; Meulenberg et al., *supra*).

The existence of apathogenic or low-pathogenic strains among isolates has also been suggested (Stevenson, *supra*). Thus, these studies suggest that the PRRSV isolates in

North America and in Europe are antigenically and genetically heterogeneous, and that different genotypes or serotypes of PRRSV exist. However, prior to the present invention, the role of antigenic and genetic variation in the pathogenesis of PRRSV was not entirely clear.

The occurrence of PRRS in the U.S. has adversely affected the pig farming industry. Almost half of swine herds in swine-producing states in the U.S. are seropositive for PRRSV (*Animal Pharm.*, 264:11 (11/11/92)). In Canada, PRRS has been characterized by anorexia and pyrexia in sows lasting up to 2 weeks, late-term abortions, increased stillbirth rates, weak-born pigs and neonatal deaths preceded by rapid abdominal breathing and diarrhea. Work on the isolation of the virus causing PRRS, on a method of diagnosing PRRS infection, and on the development of a vaccine against the PRRS virus has been published (see Canadian Patent Publication No. 2,076,744; PCT International Patent Publication No. WO 93/03760; PCT International Patent Publication No. WO 93/06211; and PCT International Patent Publication No. WO 93/07898).

There is also variability in the virulence of PRRSV in herds. Recently, a more virulent form of PRRS has been occurring with increased incidence in 3-8 week old pigs in the midwestern United States. Typically, healthy 3-5 week old pigs are weaned and become sick 5-7 days later. Routine virus identification methods on tissues from affected pigs have shown that swine influenza virus (SIV), pseudorabies virus (PRV), and Mycoplasma hyopneumoniae are not associated with this new form of PRRS. Originally termed proliferative interstitial pneumonia (PIP; see U.S. patent application Serial No. 07/969,071), this disease has been very recently linked with PRRS, and the virus has been informally named the "Iowa strain" of PRRSV (see U.S. patent application Serial No. 08/131,625).

Pessimism and skepticism has been expressed in the art concerning the development of effective vaccines against these porcine viruses (*The Veterinary Record*, October 26, 1991). A belief that human influenza vaccine may afford some protection against the effects of PRRS and PNP exists (*The Veterinary Record*, July 6, 1991).

Viral envelope proteins are known to be highly variable in many coronaviruses, such as feline infectious peritonitis virus and mouse hepatitis virus (Dalziel et al: Site-specific alteration of murine hepatitis virus type 4 peplomer glycoprotein E2 results in reduced neurovirulence. *J. Virol.*, 59:464-471 (1986); Fleming et al: Pathogenicity of antigenic variants of murine coronavirus JHM selected with monoclonal antibodies. *J. Virol.*, 58:869-875 (1986); Fiscus et al: Antigenic comparison of the feline coronavirus isolates; Evidence for markedly different peplomer glycoproteins. *J. Virol.*, 61:2607-2613 (1987); Parker et al: Sequence analysis reveals extensive polymorphism and evidence of deletions within the E2 glycoprotein gene of several strains of murine hepatitis virus. *Virology*, 173:664-673 (1989)).

For example, a deletion or a mutation in the major envelope protein in coronaviruses can alter tissue tropism and *in vivo* pathogenicity. A mutation in a monoclonal antibody-resistant mutant of MHV has resulted in loss of its neurovirulence for mice (Fleming et al, 1986 *supra*). Porcine respiratory coronavirus (PRCV) is believed to be a deletion mutant of transmissible gastroenteritis virus (TGEV) in swine. The deletion in the PRCV genome may be in the 5'-end of the spike (S) gene of TGEV (Halbur et al, An overview of porcine viral respiratory disease. *Proc. Central Veterinary Conference*, pp. 50-59 (1993); Laude et al, Porcine respiratory coronavirus: Molecular features and virus-host interactions. *Vet. R s.*, 24:125-150 (1993); Vaughn et al, Isolation and characterization of thr e

porcine respiratory coronavirus isolates with varying sizes of deletions. *J. Clin. Micro.*, 32:1809-1812 (1994)).

PRCV has a selective tropism for the respiratory tract and does not replicate in the gastrointestinal tract (Rasschaert et al, Porcine respiratory coronavirus differs from transmissible gastroenteritis virus by a few genomic deletions. *J. Gen. Virol.*, 71:2599-2607 (1990); Laude et al, 1993 *supra*). In contrast, TGEV has a tropism for both respiratory and gastrointestinal tracts (Laude et al, 1993 *supra*).

Variation in antigenic and genetic relatedness among LDV isolates of varying pathogenicity is also known (Kuo et al, Lactate-dehydrogenase-elevating virus (LDV): subgenomic mRNAs, mRNA leader and comparison of 3'-terminal sequences of two LDV isolates. *Virus Res.*, 23:55-72 (1992); Plagemann, LDV, EAV, and SHFV: A new group of positive stranded RNA viruses. *Proc. Am. Assoc. Swine Practitioners*, 4:8-15 (1992); Chen et al, Sequences of 3' end of genome and of 5' end of open reading frame 1a of lactate dehydrogenase-elevating virus and common junction motifs between 5' leader and bodies of seven subgenomic mRNAs. *J. Gen. Virol.*, 74:643-660 (1993)).

However, the present invention provides the first insight into the relationships between the open reading frames of the PRRSV genome and their corresponding effects on virulence and replication.

Further, a diagnosis of porcine reproductive and respiratory syndrome (PRRS) relies on compiling information from the clinical history of the herd, serology, pathology, and ultimately on isolation of the PRRS virus (PRRSV). Three excellent references reviewing diagnosis of PRRSV have been published in the last year (Van Alstine et al, "Diagnosis of porcine reproductive and respiratory syndrome," *Swine Health and Production*, Vol. 1, No. 4 (1993), p. 24-28; Christianson et al, "Porcine reproductiv

and respiratory syndrome: A review." *Swine Health and Production*, Vol. 1, No. 2 (1994), pp. 10-28 and Goyal, "Porcine reproductive and respiratory syndrome," *J. Vet. Diagn. Invest.* 5:656-664 (1993)). PRRSV has also recently been shown to replicate in pulmonary alveolar macrophages by gold colloid immunohistochemistry (Magar et al (1993): Immunohistochemical detection of porcine reproductive and respiratory syndrome virus using colloidal gold. *Can. J. Vet. Res.*, 57:300-304).

Clinical signs vary widely between farms, and thus, are not the most reliable evidence of a definitive diagnosis, except in the case of a severe acute outbreak in naive herds which experience abortion storms, increased numbers of stillborn pigs, and severe neonatal and nursery pig pneumonia. Presently, the most common clinical presentation is pneumonia and miscellaneous bacterial problems in 3-10 week old pigs. However, many PRRSV-positive herds have no apparent reproductive or respiratory problems.

Some herds evidence devastating reproductive failure, characterized by third-trimester abortions, stillborn pigs and weak-born pigs. Many of these herds also experience severe neonatal respiratory disease. Respiratory disease induced by PRRSV in 4-10 week-old pigs is also common and can be quite severe (Halbur et al, *Viral contributions to the porcine respiratory disease complex*. *Proc. Am. Assoc. Swine Pract.* (1993), pp. 343-350). Clinical PRRSV outbreaks are frequently followed by bacterial pneumonia, septicemia, or enteritis. Thus, it has been difficult to obtain an acceptably rapid and reliable diagnosis of infection by PRRSV, prior to the present invention.

The pig farming industry has been and will continue to be adversely affected by these porcine reproductive and respiratory diseases and new variants thereof, as they appear. PRRSV is a pathogen of swine that causes economic

losses from reproductive and respiratory diseases. Economic losses from PRRS occur from loss of pigs from abortions, stillborn pigs, repeat breeding, pre-weaning and postweaning mortality, reduced feed conversion efficiency, increased drug and labor cost and have been estimated to cost approximately \$236 per sow in addition to loss of profits (Polson et al., Financial implications of mystery swine disease (MSD), *Proc. Mystery Swine Disease Committee Meeting*, Denver, Co., 1990, pp. 8-28). This represents a loss of \$23,600 for a 100 sow herd or \$236,000 for a 1000 sow herd.

PRRSV causes additional losses from pneumonia in nursery pigs. However, the exact economic losses from PRRSV-associated pneumonia are not known. PRRSV is an important cause of pneumonia in nursery and weaned pigs. Reproductive disease was the predominant clinical outcome of PRRSV infections during the past few years. Respiratory disease has now become the main problem associated with PRRSV.

Surprisingly, the market for animal vaccines in the U.S. and worldwide is larger than the market for human vaccines. Thus, there exists an economic incentive to develop new veterinary vaccines, in addition to the substantial public health benefit which is derived from protecting farm animals from disease.

Disclosure of the Invention

Accordingly, one object of the present invention is to provide a polynucleic acid isolated from a porcine reproductive and respiratory virus (PRRSV).

It is a further object of the present invention to provide an isolated polynucleic acid which encodes a PRRSV prot in.

It is a further object of the present invention to provide a PRRSV protein, either isolated from a PRRSV or encoded by a PRRSV polynucleic acid.

It is a further object of the present invention to provide a protein- or polynucleic acid-based vaccine which protects a pig against PRRS.

It is a further object of the present invention to provide a method of raising an effective immunological response against a PRRSV using the vaccine.

It is a further object of the present invention to provide a method of producing a protein- or polynucleic acid-based vaccine which protects a pig against a PRRSV infection.

It is a further object of the present invention to provide a method of treating a pig infected by or exposed to a PRRSV.

It is a further object of the present invention to provide a method of detecting PRRSV.

It is a further object of the present invention to provide an immunoperoxidase diagnostic assay for detection of PRRSV antigen in porcine tissues.

It is a further object of the present invention to provide an antibody which immunologically binds to a PRRSV protein or to an antigenic region of such a protein.

It is a further object of the present invention to provide an antibody which immunologically binds to a protein- or polynucleic acid-based vaccine which protects a pig against a PRRSV.

It is a further object of the present invention to provide a method of treating a pig exposed to or infected by a PRRSV.

It is a further object of the present invention to provide a method of detecting and a diagnostic kit for assaying a PRRSV.

It is a further object of the present invention to provide the above objects, where the PRRS virus is the Iowa strain of PRRSV.

These and other objects which will become apparent during the following description of the preferred embodiments, have been provided by at least one purified polypeptide selected from the group consisting of proteins encoded by one or more open reading frames (ORF's) of an Iowa strain of porcine reproductive and respiratory virus (PRRSV), proteins at least 80% but less than 100% homologous with those encoded by one or more of ORF 2, ORF 3, ORF 4 and ORF 5 of an Iowa strain of PRRSV, proteins at least 97% but less than 100% homologous with proteins encoded by one or both of ORF 6 and ORF 7 of an Iowa strain of PRRSV, antigenic regions of said proteins which are at least 5 amino acids in length and which effectively stimulate immunological protection in a porcine host against a subsequent challenge with a PRRSV isolate, and combinations thereof; an isolated polynucleic acid which encodes such a polypeptide or polypeptides; a vaccine comprising an effective amount of such a polynucleotide or polypeptide(s); antibodies which specifically bind to such a polynucleotide or polypeptide; methods of producing the same; and methods of raising an effective immunological response against a PRRSV, treating a pig exposed to or infected by a PRRSV, and detecting a PRRSV using the same.

Brief Description of the Drawings

Figure 1 is a flowchart outlining a procedure for producing a subunit vaccine;

Figure 2 is a flowchart outlining a procedure for producing a genetically engineered vaccine;

Figure 3 shows a general schematic procedure for the construction of a cDNA λ library as described by the manufacturer (Stratagene);

Figure 4 shows a general schematic procedure for identifying authentic clones of the PRRS virus isolate ISU-12 (VR 2385) by differential hybridization (modified from "Recombinant DNA," 2nd ed., Watson, J.D., et al., eds. (1992), p. 110);

Figure 5 is a Northern blot showing the VR 2385 subgenomic mRNA species, denatured with 6 M glyoxal and DMSO, and separated on a 1.5% agarose gel;

Figure 6 shows the λ cDNA clones used to obtain the 3'-terminal nucleotide sequence of VR 2385;

Figure 7 shows the 2062-bp 3'-terminal sequence (SEQ ID NO:13) and the amino acid sequences encoded by ORF's 5, 6 and 7 (SEQ ID NOS:15, 17 and 19, respectively) of VR 2385;

Figure 8 compares the ORF-5 regions of the genomes of VR 2385 and Lelystad virus;

Figure 9 compares the ORF-6 regions of the genomes of VR 2385 and Lelystad virus;

Figure 10 compares the ORF-7 regions of the genomes of VR 2385 and Lelystad virus;

Figure 11 compares the 3'-nontranslational regions of the genomes of VR 2385 and Lelystad virus;

Figure 12 shows a cytopathic effect in HI-FIVE cells infected with a recombinant baculovirus containing the VR 2385 ORF-7 gene (Baculo.PRRSV.7);

Figure 13 shows HI-FIVE cells infected with a recombinant baculovirus containing the VR 2385 ORF-6 gene, stained with swine antisera to VR 2385, followed by fluorescein-conjugated anti-swine IgG;

Figure 14 shows HI-FIVE cells infected with a recombinant baculovirus containing the VR 2385 ORF-7 gene, respectively, stained with swine antisera to VR 2385, followed by fluorescein-conjugated anti-swine IgG;

Figure 15 shows a band of expected size for the VR 2385 ORF-6 product, detected by a radioimmunoprecipitation technique (see Experiment II(B) below);

Figure 16 shows a band of expected size for the VR 2385 ORF-7 product, detected by a radioimmunoprecipitation technique (see Experiment II(B) below);

Figure 17 compares the ORF 6 and ORF 7 nucleotide sequences of six U.S. PRRSV isolates and of LV, in which the VR 2385 nucleotide sequence is shown first, and in subsequent sequences, only those nucleotides which are different are indicated;

Figures 18(A)-(B) show the alignment of amino acid sequences of the putative M (Fig. 18(A)) and N (Fig. 18(B)) genes of the proposed arterivirus group, performed with a GENEWORKS program (IntelliGenetics, Inc.);

Figures 19(A)-(B) show phylogenetic trees based on the amino acid sequences of the putative M (Fig. 19(A)) and N genes (Fig. 19(B)) for the proposed arterivirus group;

Figure 20 shows the nucleotide sequence of a region of the genome of PRRSV isolate VR 2385 containing ORF's 2, 3 and 4;

Figures 21(A)-(C) compare the nucleotide sequences of ORF 2, ORF 3 and ORF 4 of PRRSV VR 2385 with the corresponding ORF's of Lelystad virus (LV);

Figures 22(A)-(C) show alignments of the predicted amino acid sequences encoded by ORF's 2, 3 and 4 of PRRSV VR 2385 and LV;

Figure 23 shows an immunohistochemical stain of a lung tissue sample taken from a pig infected 9 days previously with PRRSV, in which positive ABC staining with hematoxylin counterstain is observed within the cytoplasm of macrophages and sloughed cells in the alveolar spaces;

Figure 24 shows an immunohistochemical stain of a lung tissue sample taken from a pig infected 4 days previously with PRRSV, in which positive ABC staining with hematoxylin

counterstain is demonstrated within cellular debris in terminal airway lumina;

Figure 25 shows a heart from a pig infected 9 days previously with PRRSV, in which positive staining is demonstrated within endothelial cells (arrow) and isolated macrophages by the present streptavidin-biotin complex method (with hematoxylin counterstain); the bar indicates a length of 21 microns;

Figure 26 shows a tonsil from a pig infected 9 days previously with PRRSV, in which positive staining cells (arrow heads) are demonstrated within follicles and in the crypt epithelium by the present streptavidin-biotin complex method (with hematoxylin counterstain); the bar indicates a length of 86 microns;

Figure 27 shows a lymph node from a pig infected 9 days previously with PRRSV, in which positive staining is demonstrated within follicles by the present streptavidin-biotin complex method (with hematoxylin counterstain), and positive cells (arrows) resemble macrophages or dendritic cells; the bar indicates a length of 21 microns;

Figures 28(A)-(C) are photomicrographs of lungs from pig inoculated with (A) culture fluid from an uninfected cell line, (B) culture fluid from a cell line infected with a low virulence PRRSV isolate (the lungs show PRRS-A type lesions), and (C) culture fluid from a cell line infected with a high virulence PRRSV isolate (the lungs show PRRS-B type lesions);

Figures 29(A)-(B) illustrate immunohistochemical staining with anti-PRRSV monoclonal antibody of a lung from a pig infected 9 days previously with PRRSV; and

Figures 30(A)-(B) show Northern blots of PRRSV isolates VR 2385pp (designated as "12"), VR 2429 (ISU-22, designated as "22"), VR 2430, designated as "55"), ISU-79 (designated as "79"), ISU-1894 (designated as "1894"), and VR 2431, designated as "3927").

Best Mode for Carrying Out the Invention

In the present invention, a "porcine reproductive and respiratory syndrome virus" or "PRRSV" refers to a virus which causes the diseases PRRS, PEARs, SIRS, MSD and/or PIP (the term "PIP" now appears to be disfavored), including the Iowa strain of PRRSV, other strains of PRRSV found in the United States (e.g., VR 2332), strains of PRRSV found in Canada (e.g., IAF-exp91), strains of PRRSV found in Europe (e.g., Lelystad virus, PRRSV-10), and closely-related variants of these viruses which may have appeared and which will appear in the future.

The present vaccine is effective if it protects a pig against infection by a porcine reproductive and respiratory syndrome virus (PRRSV). A vaccine protects a pig against infection by a PRRSV if, after administration of the vaccine to one or more unaffected pigs, a subsequent challenge with a biologically pure virus isolate (e.g., VR 2385, VR 2386, or other virus isolate described below) results in a lessened severity of any gross or histopathological changes (e.g., lesions in the lung) and/or of symptoms of the disease, as compared to those changes or symptoms typically caused by the isolate in similar pigs which are unprotected (i.e., relative to an appropriate control). More particularly, the present vaccine may be shown to be effective by administering the vaccine to one or more suitable pigs in need thereof, then after an appropriate length of time (e.g., 1-4 weeks), challenging with a large sample (10^{3-7} TCID₅₀) of a biologically pure PRRSV isolate. A blood sample is then drawn from the challenged pig after about one week, and an attempt to isolate the virus from the blood sample is then performed (e.g., see the virus isolation procedure exemplified in Experiment VIII below). Isolation of the virus is an indication that the vaccine may not be

effective, and failure to isolate the virus is an indication that the vaccine may be effective.

Thus, the effectiveness of the present vaccine may be evaluated quantitatively (i.e., a decrease in the percentage of consolidated lung tissue as compared to an appropriate control group) or qualitatively (e.g., isolation of PRRSV from blood, detection of PRRSV antigen in a lung, tonsil or lymph node tissue sample by an immunoperoxidase assay method [described below], etc.). The symptoms of the porcine reproductive and respiratory disease may be evaluated quantitatively (e.g., temperature/fever), semi-quantitatively (e.g., severity of respiratory distress [explained in detail below], or qualitatively (e.g., the presence or absence of one or more symptoms or a reduction in severity of one or more symptoms, such as cyanosis, pneumonia, heart and/or brain lesions, etc.).

An unaffected pig is a pig which has either not been exposed to a porcine reproductive and respiratory disease infectious agent, or which has been exposed to a porcine reproductive and respiratory disease infectious agent but is not showing symptoms of the disease. An affected pig is one which shows symptoms of PRRS or from which PRRSV can be isolated.

The clinical signs or symptoms of PRRS may include lethargy, respiratory distress, "thumping" (forced expiration), fevers, roughened haircoats, sneezing, coughing, eye edema and occasionally conjunctivitis. Lesions may include gross and/or microscopic lung lesions, myocarditis, lymphadenitis, encephalitis and rhinitis. The infectious agent may be a single virus, or may be combined with one or more additional infectious agents (e.g., other viruses or bacteria). In addition, less virulent and non-virulent forms of the PRRSV and of Iowa strain have been found, which may cause either a subset of the above symptoms or no symptoms at all. Less virulent and non-

virulent forms of PRRSV can be used according to the present invention to provide protection against porcine reproductive and respiratory diseases nonetheless.

Histological lesions in the various porcine diseases are different. Table I below compares physiological observations and pathology of the lesions associated with a number of diseases caused by porcine viruses:

TABLE I

Swine Viral Pneumonia Comparative Pathology

Lesion	PRRS(p)	PRRS(o)	SIV	PNP	PRCV	PPMV	Iowa
Type II	+	+++	+	+++	++	++	++++
Inter. thickening	++++	+	+	+	++	++	+
Alveolar exudate	+	+++	++	++	++	++	+++
Airway necrosis	-	-	++++	++++	+++	+	-
Syncytia	-	++	+/-	++	+	+	+++
Encephalitis	+	+++	-	-	-	++	+
Myocarditis	+/-	++	-	-	-	-	+++

wherein "PRRS(p)" represents the published pathology of the PRRS virus, "PRRS(o)" represents the pathology of PRRS virus observed by the present Inventors, "SIV" represents swine influenza A virus, "PRCV" represents porcine respiratory coronavirus, "PPMV" represents porcine paramyxovirus, "Iowa" refers to the strain of PRRSV discovered by the present Inventors, "Type II" refers to Type II pneumocytes (which proliferate in infected pigs), "Inter." refers to interstitial septal infiltration by mononuclear cells, "Airway necrosis" refers to necrosis in terminal airways, and the symbols (-) and (+) through (++++), refer to a comparative severity scal as follows:

-21-

(-):negative (not observed)

(+):mild (just above the threshold of observation)

(++):moderate

(+++):severe

(++++):most severe

A "porcine reproductive and respiratory virus" or "PRRSV" causes a porcine reproductive and respiratory disease defined by one or more of the clinical signs, symptoms, lesions and histopathology as described above, and is characterized as being an enveloped RNA arterivirus, having a size of from 50 to 80 nm in diameter and from 250 to 400 nm in length. "North American strains of PRRSV" refer to those strains of PRRSV which are native to North America. "U.S. strains of PRRSV" refer to strains of PRRSV native to the U.S., and "European strains of PRRSV" refer to strains native to Europe, such as Lelystad virus (deposited by the CDI [Lelystad, Netherlands] in the depository at the Institut Pasteur, Paris, France, under the deposit number I-1102; see International Patent Publication No. WO 92/21375, published on December 10, 1992).

The "Iowa strain" of PRRSV refers to (a) those strains of PRRSV isolated by the presented Inventors, (b) those strains having at least a 97% sequence identity (or homology) in the seventh open reading frame (ORF 7) with at least one of VR 2385, VR 2430 and VR 2431; (c) strains which, after no more than 5 passages, grow to a titer of at least 10^4 TCID₅₀ in CRL 11171 cells, MA-104 cells or PSP-36 cells, (d) those strains having at least 80% and preferably at least 90% homology with one or more of ORF's 2-5 of VR 2385, and (e) those strains which cause a greater percentage consolidation of lung tissue than Lelystad virus

(e.g., at 10 days post-infection, infected pigs exhibit at least 20% and preferably at least 40% lung consolidation). Preferably, the Iowa strain of PRRSV is characterized by at least two of the above characteristics (a)-(e).

The present invention is primarily concerned with polynucleic acids (segments of genomic RNA and/or DNA, mRNA, cDNA, etc.) isolated from or corresponding to a porcine reproductive and respiratory syndrome virus (PRRSV), proteins encoded by the DNA, methods of producing the polynucleic acids and proteins, vaccines which protect pigs from a PRRSV, a method of protecting a pig from a PRRSV using the vaccine, a method of producing the vaccine, a method of treating a pig infected by or exposed to a PRRSV, and a method of detecting a PRRSV. More particularly, the present invention is concerned with a vaccine which protects pigs from North American strains of PRRSV, a method of producing and administering the vaccine, and polynucleic acids and proteins obtained from an Iowa strain of PRRSV. However, it is believed that the information learned in the course of developing the present invention will be useful in developing vaccines and methods of protecting pigs against any and/or all strains of porcine reproductive and respiratory syndrome. Therefore, the present invention is not necessarily limited to polynucleic acids, proteins, vaccines and methods related to the Iowa strain of PRRSV virus (PRRSV).

The phrase "polynucleic acid" refers to RNA or DNA, as well as mRNA and cDNA corresponding to or complementary to the RNA or DNA isolated from the virus or infectious agent. An "ORF" refers to an open reading frame, or polypeptide-encoding segment, isolated from a viral genome, including the PRRSV genome. In the present polynucleic acid, an ORF can be included in part (as a fragment) or in whole, and can overlap with the 5'- or 3'-sequence of an adjacent ORF (see Figs. 7 and 21, and Experiments I and IV below). A

"polynucleotide" is equivalent to a polynucleic acid, but may define a distinct molecule or group of molecules (e.g., as a subset of a group of polynucleic acids).

Referring now to Figures 1-2, flowcharts of procedures are provided for preparing types of vaccines encompassed by the present invention. The flowcharts of Figures 1-2 are provided as exemplary methods of producing the present vaccines, and are not intended to limit the present invention in any manner.

The first step in each procedure detailed in Figures 1-2 is to identify a cell line susceptible to infection with a porcine reproductive and respiratory virus or infectious agent. (To simplify the discussion concerning preparation of the vaccine, the term "virus" refers to a virus and/or other infectious agent associated with a porcine reproductive and respiratory disease.) A master cell stock (MCS) of the susceptible host cell is then prepared. The susceptible host cells continue to be passaged beyond MCS. Working cell stock (WCS) is prepared from cell passages between MCS and MCS+n.

A master seed virus is propagated on the susceptible host cell line, between MCS and MCS+n, preferably on WCS. The raw virus is isolated by methods known in the art from appropriate, preferably homogenized, tissue samples taken from infected pigs exhibiting disease symptoms corresponding to those caused by the virus of interest. A suitable host cell, preferably a sample of the WCS, is infected with the raw virus, then cultured. Vaccine virus is subsequently isolated and plaque-purified from the infected, cultured host cell by methods known in the art. Preferably, the virus to be used to prepare the vaccine is plaque-purified three times.

Master seed virus (MSV) is then prepared from the plaque-purified virus by methods known in the art. The MSV(X) is then passaged in WCS at least four times through

MSV(X+1), MSV(X+2), MSV(X+3) and MSV(X+4) virus passages. The MSV(X+4) is considered to be the working seed virus. Preferably, the virus passage to be used in the pig studies and vaccine product of the present invention is MSV(X+5), the product of the fifth passage.

In conjunction with the working cell stock, the working seed virus is cultured by known methods in sufficient amounts to prepare a prototype vaccine, preferably MSV(X+5). The present prototype vaccines may be of any type suitable for use in the veterinary medicine field. The primary types of vaccines on which the present invention focuses include a subunit vaccine (Figure 1) and a genetically engineered vaccine (Figure 2). However, other types of vaccines recognized in the field of veterinary vaccines, including live, modified live, attenuated and killed virus vaccines, are also acceptable. A killed vaccine may be rendered inactive through chemical treatment or heat, etc., in a manner known to the artisan of ordinary skill.

An attenuated virus may be obtained by repeating serial passage of the virus in a suitable host cell a sufficient number of times to obtain an essentially non-virulent virus. For example, a PRRSV may be serially passaged from 1 to 20 times (or more, if desired), in order to render it sufficiently attenuated for use as an attenuated vaccine. MSV(X+5) may be such an attenuated vaccine.

In the procedures outlined by each of Figures 1-2, following preparation of a prototype vaccine, pig challenge models and clinical assays are conducted by methods known in the art. For example, before performing actual vaccination/challenge studies, the disease to be prevented and/or treated must be defined in terms of its symptoms, clinical assay results, conditions, etc. As described herein, the Iowa strain of PRRSV has been defined in terms

of its histopathology and the clinical symptoms which it causes. Clinical analyses of the Iowa strain of PRRSV are described in detail in the Experiments below.

One then administers a prototype vaccine to a pig, then exposes the pig to the virus which causes the disease. This is known as "challenging" the pig and its immunological system. After observing the response of the challenged pig to exposure to the virus or infectious agent and analyzing the ability of the prototype vaccine to protect the pig, efficacy studies are then performed by conventional, known methods. A potency assay is then developed in a separate procedure by methods known in the art, and prelicensing serials are then produced.

Prior to preparation of the prototype subunit vaccine (Figure 1), the protective or antigenic components of the vaccine virus should be identified. Such protective or antigenic components include certain amino acid segments or fragments of the viral proteins (preferably coat proteins) which raise a particularly strong protective or immunological response in pigs; such antigenic protein fragments fused to non-PRRSV proteins which act as a carrier and/or adjuvant; single or multiple viral coat proteins themselves, oligomers thereof, and higher-order associations of the viral coat proteins which form virus substructures or identifiable parts or units of such substructures; oligoglycosides, glycolipids or glycoproteins present on or near the surface of the virus or in viral substructures such as the nucleocapsid; lipoproteins or lipid groups associated with the virus, etc.

Antigenic amino acid segments or fragments are preferably at least 5 amino acids in length, particularly preferably at least 10 amino acids in length, and can be up to but not including the entire length of the native protein. In the present invention, the binding affinity

(or binding constant or association constant) of an antigenic fragment is preferably at least 1% and more preferably at least 10% of the binding affinity of the corresponding full-length protein (i.e., which is encoded by the same ORF) to a monoclonal antibody which specifically binds the full-length protein. The monoclonal antibody which specifically binds to the full-length protein encoded by an ORF of a PRRSV is preferably deposited under the Budapest Treaty at an acceptable depository, or is sequenced or otherwise characterized in terms of its physicochemical properties (e.g., antibody type [IgG, IgM, etc.], molecular weight, number of heavy and light chains, binding affinities to one or more known or sequenced proteins [e.g., selected from SEQ ID NOS:15, 17, 19, 21, 24, 26, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 67, 69, 71, 73, 75 and 77], etc.).

Antigenic fragments of viral proteins (e.g., those encoded by one or more of ORF's 2-6 of a PRRSV virus) are identified by methods known in the art. For example, one can prepare polynucleic acids having a truncated ORF encoding a polypeptide with a predetermined number of amino acid residues deleted from the N-terminus, C-terminus, or both. The truncated ORF can be expressed *in vitro* or *in vivo* in accordance with known methods, and the corresponding truncated polypeptide can then be isolated in accordance with known methods. The immunoprotective properties of the polypeptides may be measured directly (e.g., *in vivo*). Alternatively, the antigenic region(s) of the full-length polypeptide can be determined indirectly by screening a series of truncated polypeptides against, for example, suitably deposited or characterized monoclonal antibodies. (If the alternative, indirect method is performed, the failure of a truncated polypeptide to bind to a neutralizing monoclonal antibody is a strong indication that the portion of the full-length polypeptide

deleted in the truncated polypeptide contains an antigenic fragment.) Once identified, the antigenic or immunoprotective portion(s) (the "subunit(s)") of the viral proteins or of the virus itself may be subsequently cloned and/or purified in accordance with known methods. (The viral/bacterial inactivation and subunit purification protocols recited in Fig. 1 are optional.)

Genetically engineered vaccines (Figure 2) begin with a modification of the general procedure used for preparation of the other vaccines. After plaque-purification, the PRRS virus may be isolated from a suitable tissue homogenate by methods known in the art, preferably by conventional cell culture methods using PSP-36, ATCC CRL 11171 or macrophage cells as hosts.

The RNA is extracted from the biologically pure virus by a known method, preferably by the guanidine isothiocyanate method using a commercially available RNA isolation kit (for example, the kit available from Stratagene, La Jolla, California), and purified by one or more known methods, preferably by ultracentrifugation in a CsCl gradient. Messenger RNA may be further purified or enriched by oligo (dT)-cellulose column chromatography.

The viral genome is then cloned into a suitable host by methods known in the art (see Maniatis et al, "Molecular Cloning: A Laboratory Manual," Cold Spring Harbor Laboratory (1989), Cold Spring Harbor, Massachusetts). The virus genome is then analyzed to determine essential regions of the genome for producing antigenic portions of the virus. Thereafter, the procedure for producing a genetically engineered vaccine is essentially the same as for a modified live vaccine, an inactivated vaccine or a subunit vaccine (see Figure 1 of the present application and Figures 1-3 of U.S. application Serial No. 08/131,625). During pre licensing serials, expression of the cloned, recombinant subunit of a subunit vaccine may be optimized

by methods known to those in the art (see, for example, relevant sections of Maniatis et al, cited above).

The present vaccine protects pigs against a virus or infectious agent which causes a porcine reproductive and respiratory disease. Preferably, the present vaccine protects pigs against infection by PRRSV. However, the present vaccine is also expected to protect a pig against infection by closely related variants of various strains of PRRSV as well.

Subunit virus vaccines may also be prepared from semi-purified virus subunits by the methods described above in the discussion of Figure 1. For example, hemagglutinin isolated from influenza virus and neuraminidase surface antigens isolated from influenza virus have been prepared, and shown to be less toxic than the whole virus. Subunit vaccines can also be prepared from highly purified subunits of the virus. An example in humans is the 22-nm surface antigen of human hepatitis B virus. Human herpes simplex virus subunits and many other examples of subunit vaccines for use in humans are known. Thus, methods of preparing purified subunit vaccines from PRRSV cultured in a suitable host cell may be applicable to the present subunit vaccine.

Attenuated virus vaccines can be found in nature and may have naturally-occurring gene deletions (see Experiments VIII and IX below). Alternatively, attenuated vaccines may be prepared by a variety of known methods, such as serial passage (e.g., 5-25 times) in cell cultures or tissue cultures. However, the attenuated virus vaccines preferred in the present invention are those attenuated by recombinant gene deletions or gene mutations (as described above).

Genetically engineered vaccines are produced by techniques known to those in the art. Such techniques include those using recombinant DNA and those using live viruses. For example, certain virus genes can be

identified which code for proteins responsible for inducing a stronger immune or protective response in pigs. Such identified genes can be cloned into protein expression vectors, such (but not limited to) as the baculovirus vector (see, for example, O'Reilly et al, "Baculovirus Expression Vectors: A Lab Manual," Freeman & Co. (1992)). The expression vector containing the gene encoding the immunogenic virus protein can be used to infect appropriate host cells. The host cells are cultured, thus expressing the desired vaccine proteins, which can be purified to a desired extent, then used to protect the pigs from a reproductive and respiratory disease.

Genetically engineered proteins may be expressed, for example, in insect cells, yeast cells or mammalian cells. The genetically engineered proteins, which may be purified and/or isolated by conventional methods, can be directly inoculated into animals to confer protection against porcine reproductive and respiratory diseases. One or more envelope proteins from a PRRSV (i.e., those encoded by ORF's 2-6) or antigenic portions thereof may be used in a vaccine to induce neutralizing antibodies. Nucleoproteins from a PRRSV may be used in a vaccine to induce cellular immunity.

Preferably, the present invention transforms an insect cell line (HI-FIVE) with a transfer vector containing polynucleic acids obtained from the Iowa strain of PRRSV. Preferably, the present transfer vector comprises linearized baculovirus DNA and a plasmid containing one or more polynucleic acids obtained from the Iowa strain of PRRSV. The host cell line may be co-transfected with the linearized baculovirus DNA and a plasmid, so that a recombinant baculovirus is made. Particularly preferably, the present polynucleic acid encodes one or more proteins of the Iowa strain of PRRSV.

Alternatively, RNA or DNA from a PRRSV encoding one or more viral proteins (e.g., envelope and/or nucleoproteins) can be inserted into live vectors, such as a poxvirus or an adenovirus, and used as a vaccine.

Thus, the present invention further concerns a purified preparation of a polynucleic acid isolated from the genome of a PRRS virus, preferably a polynucleic acid isolated from the genome of the Iowa strain of PRRSV. The present polynucleic acid has utility (or usefulness) in the production of the present vaccine, in screening or identifying infected or exposed animals, in identifying related viruses and/or infectious agents, and as a vector for transforming cells and/or immunizing animals (e.g., pigs) with heterologous genes.

In the Experiments described hereinbelow, the isolation, cloning and sequencing of ORF's 2-7 of plaque-purified PRRSV isolate ISU-12 (deposited on October 30, 1992, in the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A., under the accession numbers VR 2385 [3 x plaque-purified] and VR 2386 [non-plaque-purified]) and ORF's 6-7 of PRRSV isolates ISU-22, ISU-55 and ISU-3927 (deposited on September 29, 1993, in the American Type Culture Collection under the accession numbers VR 2429, VR 2430 and VR 2431, respectively), ISU-79 and ISU-1894 (deposited on August 31, 1994, in the American Type Culture Collection under the accession numbers VR 2474 and VR 2475, respectively) are described in detail. However, the techniques used to isolate, clone and sequence these genes can be also applied to the isolation, cloning and sequencing of the genomic polynucleic acids of any PRRSV. Thus, the present invention is not limited to the specific sequences disclosed in the Experiments below.

For example, primers for making relatively large amounts of DNA by the polymerase chain reaction (and if

desired, for making RNA by transcription and/or protein by translation in accordance with known *in vivo* or *in vitro* methods) can be designed on the basis of sequence information where more than one sequence obtained from a PRRSV genome has been determined (e.g., ORF's 2-5 of VR 2385 and Lelystad virus, or ORF's 6-7 of VR 2385, VR 2429, VR 2430, ISU-79, ISU-1894, VR 2431 and Lelystad virus). A region from about 15 to 50 nucleotides in length having at least 80% and preferably at least 90% identity is selected from the determined sequences. A region where a deletion occurs in one of the sequences (e.g., of at least 5 nucleotides) can be used as the basis for preparing a selective primer for selective amplification of the polynucleic acid of one strain or type of PRRSV over another (e.g., for the differential diagnosis of North American and European PRRSV strains).

Once the genomic polynucleic acid is amplified and cloned into a suitable host by known methods, the clones can be screened with a probe designed on the basis of the sequence information disclosed herein. For example, a region of from about 50 to about 500 nucleotides in length is selected on the basis of either a high degree of identity (e.g., at least 90%) among two or more sequences (e.g., in ORF's 6-7 of the Iowa strains of PRRSV disclosed in Experiment III below), and a polynucleotide of suitable length and sequence identity can be prepared by known methods (such as automated synthesis, or restriction of a suitable fragment from a polynucleic acid containing the selected region, PCR amplification using primers which hybridize specifically to the polynucleotide, and isolation by electrophoresis). The polynucleotide may be labeled with, for example, ³²P (for radiometric identification) or biotin (for detection by fluorometry). The probe is then hybridized with the polynucleic acids of the clones and detected according to known methods.

The present Inventors have discovered that ORF 4 appears to be related to the virulence of PRRSV. For example, at least one isolate of PRRSV which shows relatively low virulence also appears to have a deletion in ORF 4 (see, for example, Experiments VIII-XI below). Accordingly, in a preferred embodiment, the present invention is concerned with a polynucleic acid obtained from a PRRSV isolate which confers immunogenic protection directly or indirectly against a subsequent challenge with a PRRSV, but in which ORF 4 is deleted or mutated to an extent which would render a PRRSV containing the polynucleic acid either low-virulent (i.e., a "low virulence" (lv) phenotype; see the explanation below) or non-virulent (a so-called "deletion mutant"). Preferably, ORF 4 is deleted or mutated to an extent which would render a PRRSV virus non-virulent. However, it may be desirable to retain regions of a PRRSV ORF 4 in the present polynucleic acid which (i) encode an antigenic, immunoprotective peptide fragment and (ii) would not confer virulence to a PRRSV virus containing the polynucleic acid.

The present invention also encompasses a PRRSV per se in which ORF 4 is deleted or mutated to an extent which renders it either low-virulent or non-virulent (e.g., VR 2431). Such a virus is useful as a vaccine or as a vector for transforming a suitable host (e.g., MA-104, PSP 36, CRL 11171, MARC-145 or porcine alveolar macrophage cells) with a heterologous gene. Preferred heterologous genes which may be expressed using the present deletion mutant may include those encoding a protein or an antigen other than a porcine reproductive and respiratory syndrome virus antigen (e.g., pseudorabies and/or swine influenza virus proteins and/or polypeptide-containing antigens, a porcine growth hormone, etc.) or a polypeptide-based adjuvant (such as those discussed below for the present vaccine composition).

It may also be desirable in certain embodiments of the present polynucleic acid which contain, for example, the 3'-terminal region of ORF 3 (e.g., from 200 to 700 nucleotides in length), at least part of which may overlap with the 5'-region of ORF 4. Similarly, where the 3'-terminal region of ORF 4 may overlap with the 5'-terminal region of ORF 5, it may be desirable to retain the 5'-region of ORF 4 which overlaps with ORF 5.

The present Inventors have also discovered that ORF 5 in the PRRSV genome appears to be related to replication of the virus in mammalian host cells capable of sustaining a culture while infected with PRRSV. Accordingly, the present invention is also concerned with polynucleic acids obtained from a PRRSV genome in which ORF 5 may be present in multiple copies (a so-called "overproduction mutant"). For example, the present polynucleic acid may contain at least two, and more preferably, from 2 to 10 copies of ORF 5 from a high-replication (hr) phenotype PRRSV isolate.

Interestingly, the PRRSV isolate ISU-12 has a surprisingly large number of potential start codons (ATG/AUG sequences) near the 5'-terminus of ORF 5, possibly indicating alternate start sites of this gene (see SEQ ID NO:13). Thus, alternate forms of the protein encoded by ORF 5 of a PRRSV isolate may exist, particularly where alternate ORF's encode a protein having a molecular weight similar to that determined experimentally (e.g., from about 150 to about 250 amino acids in length). The most likely coding region for ORF 5 of ISU-12 (SEQ ID NO:14) is indicated in Figure 7.

One can prepare deletion and overproduction mutants in accordance with known methods. For example, one can prepare a mutant polynucleic acid which contains a "silent" or deg n rate change in the sequence of a region encoding a polypeptide. By selecting and making an appropriate degenerate mutation, one can substitute a polynucleic acid

sequence recognized by a known restriction enzyme. For example, if such a silent, degenerate mutation is made at one or two of the 3'-end of ORF 3 and the 5'- and 3'-ends of ORF 4 and ORF 5, one can insert a synthetic polynucleic acid (a so-called "cassette") which may contain multiple copies of ORF 5, multiple copies of a viral envelope protein or an antigenic fragment thereof. The "cassette" may be preceded by a suitable initiation codon (ATG), and may be suitably terminated with a termination codon at the 3'-end (TAA, TAG or TGA).

Of course, an oligonucleotide sequence which does not encode a polypeptide may be inserted, or alternatively, no cassette may be inserted. By doing so, one may provide a so-called deletion mutant.

Thus, in one embodiment of the present invention, the polynucleic acid encodes one or more proteins, or antigenic regions thereof, of a PRRSV. Preferably, the present nucleic acid encodes at least one antigenic region of a PRRSV membrane (envelope) protein. More preferably, the present polynucleic acid contains at least one copy of the ORF-5 gene from a high virulence (hv) phenotype isolate of PRRSV (see the description of "hv phenotype" below) and a sufficiently long fragment, region or sequence of at least one of ORF-2, ORF-3, ORF-4, ORF-5 and/or ORF-6 from the genome of a PRRSV isolate to encode an antigenic region of the corresponding protein(s) and effectively stimulate immunological protection against a subsequent challenge with an hv phenotype PRRSV isolate. Even more preferably, at least one entire envelope protein encoded by ORF-2, ORF-3, ORF-5 and/or ORF-6 of a PRRSV is contained in the present polynucleic acid, and the present polynucleic acid excludes a sufficiently long portion of ORF 4 from an hv PRRSV to render a PRRSV containing the same either low-virulent or non-virulent. Particularly preferably, the present polynucleic acid excludes the entire region of an

hv PRRSV ORF 4 which does not overlap with the 3'-end of ORF 3 and the 5'-end of ORF 5.

Most preferably, the polynucleic acid is isolated from the genome of an isolate of the Iowa strain of PRRSV (for example, VR 2385 (3X plaque-purified ISU-12), VR 2386 (non-plaque-purified ISU-12), VR 2428 (ISU-51), VR 2429 (ISU-22), VR 2430 (ISU-55), VR 2431 (ISU-3927), ISU-79 and/or ISU-1894).

A preferred embodiment of the present invention concerns a purified preparation which may comprise, consist essentially of or consist of a polynucleic acid having a sequence of the formula (I):

5'- α - β - γ -3' (I)

wherein α encodes at least one polypeptide or antigenic fragment thereof encoded by a polynucleotide selected from the group consisting of ORF 2 and ORF 3 of an Iowa strain of PRRSV and regions thereof encoding the antigenic fragments; and β is either a covalent bond or a linking polynucleic acid which excludes a sufficiently long portion of ORF 4 from an hv PRRSV to render the hv PRRSV either low-virulent or non-virulent; and γ is at least one copy of an ORF 5 from an Iowa strain of PRRSV, preferably from a high replication (hr) phenotype.


Alternatively, the present invention may concern a purified preparation which may comprise, consist essentially of or consist of a polynucleic acid having a sequence of the formula (II):

5'- γ - δ - ϵ -3' (II)

where γ is at least one copy of an ORF 5 from an Iowa strain of PRRSV, preferably from an hv PRRSV isolate; δ is either a covalent bond or a linking polynucleic acid which

does not materially affect transcription and/or translation of the polynucleic acid; and ϵ encodes at least one polypeptide or antigenic fragment thereof encoded by a polynucleotide selected from the group consisting of ORF 6 and ORF 7 of an Iowa strain of PRRSV and regions thereof encoding the antigenic fragments; and when δ is a covalent bond, γ may have a 3'-end which excludes the region overlapping with the 5'-end of a corresponding ORF 6. Preferably, ϵ is a polynucleotide encoding at least an antigenic region of a protein encoded by an ORF 6 of an Iowa strain of PRRSV, and more preferably, encodes at least a protein encoded by an ORF 6 of an Iowa strain of PRRSV.

The present invention may also concern a purified preparation which may comprise, consist essentially of or consist of a polynucleic acid having a sequence of the formula (III):



where α , β , γ , δ and ϵ are as defined in formulas (I) and (II) above. Thus, the present polynucleic acid may be selected from the group consisting of, from 5' to 3':

(ORF 5)_n (IV)

{-(ORF 5)_n (V)

(ORF 5)_n- η (VI)

{-(ORF 5)_n- η (VII)

where:

{ is selected from the group consisting of ORF 2-, ORF 3-, ORF 4'-, ORF 2-ORF 3-, ORF 2-ORF 4'-, ORF 3-ORF 4'- and ORF 2-ORF 3-ORF 4'-; and

η is selected from the group consisting of -ORF 5', -ORF 6, -ORF 7, -ORF 5'-ORF 6, -ORF 5'-ORF 7, -ORF 6-ORF 7 and -ORF 5'-ORF 6-ORF 7;

wherein ORF 2, ORF 3, ORF 6 and ORF 7 each encode a protein encoded by the second, third, sixth and seventh open reading frames of an Iowa strain of PRRSV, respectively; ORF 4' is a region of a fourth open reading frame of an Iowa strain of PRRSV which (i) encodes an antigenic, immunoprotective peptide fragment and which (ii) does not confer virulence to a PRRSV containing the polynucleic acid; ORF 5 is a fifth open reading frame of an hv PRRSV isolate; ORF 5' is a region of a fifth open reading frame of an Iowa strain of PRRSV which (i) encodes an antigenic, immunoprotective peptide fragment and (ii) does not confer virulence to a PRRSV containing the polynucleic acid, and which may have a 3'-end which excludes the portion overlapping with the 5'-end of a corresponding ORF 6; and $n \geq 1$.

The present polynucleic acid may also comprise, consist essentially of or consist of combinations of the above sequences, either as a mixture of polynucleotides or covalently linked in either a head-to-tail (sense-antisense) or head-to-head fashion. Polynucleic acids complementary to the above sequences and combinations thereof (antisense polynucleic acid) are also encompassed by the present invention. Thus, in addition to possessing multiple or variant copies of ORF 5, the present polynucleic acid may also contain multiple or variant copies of one or more of ORF's 1-3 and 6-7 and regions of ORF's 4-5 of Iowa strain PRRSV's.

The present invention may also concern polynucleic acids comprising, consisting essentially of or consisting of the open reading frame 1a and 1b from a PRRSV isolate. Based on information regarding viruses evolutionally related to PRRSV, ORF 1a and 1b of PRRSV are believed to encode an RNA polymase. ORF 1a and 1b are translated into a single protein by frameshifting. Preferably, the

polynucleic acid from ORF 1a and 1b of a PRRSV isolate is obtained from an Iowa strain of PRRSV.

Similar to the methods described above and in the following Experiments for ORF's 2-7, one can prepare a library of recombinant clones (e.g., using *E. coli* as a host) containing suitably prepared restriction fragments of a PRRSV genome (e.g., inserted into an appropriate plasmid expressible in the host). The clones are then screened with a suitable probe (e.g., based on a conserved sequence of ORF's 2-3; see, for example, Figure 22). Positive clones can then be selected and grown to an appropriate level. The polynucleic acids can then be isolated from the positive clones in accordance with known methods. A suitable primer for PCR can then be designed and prepared as described above to amplify the desired region of the polynucleic acid. The amplified polynucleic acid can then be isolated and sequenced by known methods.

The present purified preparation may also contain a polynucleic acid selected from the group consisting of sequences having at least 97% sequence identity (or homology) with at least one ORF 7 of VR 2385, VR 2430 and/or VR 2431; and sequences having at least 80% and preferably at least 90% sequence identity (or homology) with at least one of ORF's 1-6 of VR 2385, VR 2428, VR 2429, VR 2430 and/or VR 2431. Preferably, the polynucleic acid excludes a sufficiently long region or portion of ORF 4 of the hv PRRSV isolates VR 2385, VR 2429, ISU-28, ISU-79 and/or ISU-984 to render the isolate low-virulent or non-virulent.

In the context of the present application, "homology" refers to the percentage of identical nucleotide or amino acid residues in the sequences of two or more viruses, aligned in accordance with a conventional method for determining homology (e.g., the MACVECTOR or GENEWORKS

computer programs, aligned in accordance with the procedure described in Experiment III below).

Accordingly, a further aspect of the present invention encompasses an isolated polynucleic acid at least 90% homologous to a polynucleotide which encodes a protein, polypeptide or fragment thereof encoded by ORF's 1-7 from an Iowa strain of PRRSV (e.g., SEQ ID NOS:15, 17, 19, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65 and 67).

Preferably, the present isolated polynucleic acid encodes a protein, polypeptide, or antigenic fragment thereof which is at least 10 amino acids in length and in which amino acids non-essential for antigenicity may be conservatively substituted. An amino acid residue in a protein, polypeptide, or antigenic fragment thereof is conservatively substituted if it is replaced with a member of its polarity group as defined below:

Basic amino acids:

lysine (Lys), arginine (Arg), histidine (His)

Acidic amino acids:

aspartic acid (Asp), glutamic acid (Glu), asparagine (Asn), glutamine (Gln)

Hydrophilic, nonionic amino acids:

serine (Ser), threonine (Thr), cysteine (Cys), asparagine (Asn), glutamine (Gln)

Sulfur-containing amino acids:

cysteine (Cys), methionine (Met)

Hydrophobic, aromatic amino acids:

phenylalanine (Phe), tyrosine (Tyr), tryptophan (Trp)

Hydrophobic, nonaromatic amino acids:

glycine (Gly), alanine (Ala), valine (Val), leucine (Leu), isoleucine (Ile), proline (Pro)

More particularly, the present polynucleic acid encodes one or more of the protein(s) encoded by the second, third, fourth, fifth, sixth and/or seventh open reading frames (ORF's 2-7) of the PRRSV isolate VR 2385,

VR 2386, VR 2428, VR 2429, VR 2430, VR 2431, VR 2432, ISU-79 and/or ISU-1894 (e.g., SEQ ID NOS:15, 17, 19, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63 and 65).

Relatively short segments of polynucleic acid (about 20 bp or longer) in the genome of a virus can be used to screen or identify tissue and/or biological fluid samples from infected animals, and/or to identify related viruses, by methods described herein and known to those of ordinary skill in the fields of veterinary and viral diagnostics and veterinary medicine. Accordingly, a further aspect of the present invention encompasses an isolated (and if desired, purified) polynucleic acid consisting essentially of a fragment of from 15 to 2000 bp, preferably from 18 to 1000 bp, and more preferably from 21 to 100 bp in length, derived from ORF's 2-7 of a PRRSV genome (preferably the Iowa strain of PRRSV). Particularly preferably, the present isolated polynucleic acid fragments are obtained from a terminus of one or more of ORF's 2-7 of the genome of the Iowa strain of PRRSV, and most preferably, are selected from the group consisting of SEQ ID NOS:1-12, 22 and 28-34.

The present invention also concerns a diagnostic kit for assaying a porcine reproductive and respiratory syndrome virus, comprising (a) a first primer comprising a polynucleotide having a sequence of from 10 to 50 nucleotides in length which hybridizes to a genomic polynucleic acid from an Iowa strain of porcine reproductive and respiratory syndrome virus at a temperature of from 25 to 75°C, (b) a second primer comprising a polynucleotide having a sequence of from 10 to 50 nucleotides in length, said sequence of said second primer being found in said genomic polynucleic acid from said Iowa strain of porcine reproductive and respiratory syndrome virus and being downstream from the sequence to which the first primer hybridizes, and (c) a reagent which

enables detection of an amplified polynucleic acid. Preferably, the reagent is an intercalating dye, the fluorescent properties of which change upon intercalation into double-stranded DNA.

ORF's 6 and 7 are not likely candidates for controlling virulence and replication phenotypes of PRRSV, as the nucleotide sequences of these genes are highly conserved among high virulence (hv) and low virulence (lv) isolates (see Experiment III below). However, ORF 5 in PRRSV isolates appears to be less conserved among high replication (hr) and low replication (lr) isolates. Therefore, it is believed that the presence of an ORF 5 from an hr PRRSV isolate in the present polynucleic acid will enhance the production and expression of a recombinant vaccine produced from the polynucleic acid.

Accordingly, it is preferred that the present polynucleic acid, when used for immunoprotective purposes (e.g., in the preparation of a vaccine), contain at least one copy of ORF 5 from a high-replication isolate (i.e., an isolate which grows to a titer of 10^6 - 10^7 TCID₅₀ in, for example, CRL 11171 cells; also see the discussions in Experiments VIII-XI below).

On the other hand, the lv isolate VR 2431 appears to be a deletion mutant, relative to hv isolates (see Experiments III and VIII-XI below). The deletion appears to be in ORF 4, based on Northern blot analysis. Accordingly, when used for immunoprotective purposes, the present polynucleic acid preferably does not contain a region of ORF 4 from an hv isolate responsible for its high virulence, and more preferably, excludes the region of ORF 4 which does not overlap with the adjacent ORF's 3 and 5 (where ORF 4 overlaps with the adjacent ORF's 3 and 5).

It is also known (at least for PRRSV) that neither the nucleocapsid protein nor antibodies thereto confer immunological protection against the virus (e.g., PRRSV) to

pigs. Accordingly, the present polynucleic acid, when used for immunoprotective purposes, contains one or more copies of one or more regions from ORF's 2, 3, 4, 5 and 6 of a PRRSV isolate encoding an antigenic region of the viral envelope protein, but which does not result in the symptoms or histopathological changes associated with PRRS. Preferably, this region is immunologically cross-reactive with antibodies to envelope proteins of other PRRSV isolates. Similarly, the protein encoded by the present immunoprotective polynucleic acid confers immunological protection to a pig administered a composition comprising the protein, and antibodies to this protein are immunologically cross-reactive with the envelope proteins of other PRRSV isolates. More preferably, the present immunoprotective polynucleic acid encodes the entire envelope protein of a PRRSV isolate or a protein at least 80% homologous thereto and in which non-homologous residues are conservatively substituted, or a protein at least 90% homologous thereto.

The present isolated polynucleic acid fragments can be obtained by digestion of the cDNA corresponding to (complementary to) the viral polynucleic acids with one or more appropriate restriction enzymes, can be amplified by PCR and cloned, or can be synthesized using a commercially available automated polynucleotide synthesizer.

Another embodiment of the present invention concerns one or more proteins or antigenic fragments thereof from a PRRS virus, preferably from the Iowa strain of PRRSV. As described above, an antigenic fragment of a protein from a PRRS virus (preferably from the Iowa strain of PRRSV) is at least 5 amino acids in length, particularly preferably at least 10 amino acids in length, and provides or stimulates an immunologically protective response in a pig administered a composition containing the antigenic fragment.

Methods of determining the antigenic portion of a protein are known to those of ordinary skill in the art (see the description above). In addition, one may also determine an essential antigenic fragment of a protein by first showing that the full-length protein is antigenic in a host animal (e.g., a pig). If the protein is still antigenic in the presence of an antibody which specifically binds to a particular region or sequence of the protein, then that region or sequence may be non-essential for immunoprotection. On the other hand, if the protein is no longer antigenic in the presence of an antibody which specifically binds to a particular region or sequence of the protein, then that region or sequence is considered to be essential for antigenicity.

The present invention also concerns a protein or antigenic fragment thereof encoded by one or more of the polynucleic acids defined above, and preferably by one or more of the ORF's of a PRRSV, more preferably of the Iowa strain of PRRSV. The present proteins and antigenic fragments are useful in immunizing pigs against PRRSV, in serological tests for screening pigs for exposure to or infection by PRRSV (particularly the Iowa strain of PRRSV), etc.

For example, the present protein may be selected from the group consisting of the proteins encoded by ORF's 2-7 of VR 2385, ISU-22 (VR 2429), ISU-55 (VR 2430), ISU-1894, ISU-79 and ISU-3927 (VR 2431) (e.g., SEQ ID NOS:15, 17, 19, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 67, 69 and 71); antigenic regions of at least one of the proteins of SEQ ID NOS:15, 17, 19, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 67, 69 and 71 having a length of from 5 amino acids to less than the full length of the polypeptides of SEQ ID NOS:15, 17, 19, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 67, 69 and 71; polypeptides at least 80% homologous with a protein encoded by one of the ORF's 2-5 of VR 2385 (SEQ ID

NOS:15, 67, 69 and 71); and polypeptides at least 97% homologous with a protein encoded by one of the ORF's 6-7 of VR 2385, VR 2429, VR 2430, ISU-1894, ISU-79 and VR 2431 (e.g., SEQ ID NOS:17, 19, 43, 45, 47, 49, 51, 53, 55, 57, 59 and 61). Preferably, the present protein has a sequence selected from the group consisting of SEQ ID NOS:15, 17, 19, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 67, 69 and 71; variants thereof which provide effective immunological protection to a pig administered the same and in which from 1 to 100 (preferably from 1 to 50 and more preferably from 1 to 25) deletions or conservative substitutions in the amino acid sequence exist; and antigenic fragments thereof at least 5 and preferably at least 10 amino acids in length which provide effective immunological protection to a pig administered the same.

More preferably, the present protein variant or protein fragment has a binding affinity (or association constant) of at least 1% and preferably at least 10% of the binding affinity of the corresponding full-length, naturally-occurring protein to a monoclonal antibody which specifically binds to the full-length, naturally-occurring protein (i.e., the protein encoded by a PRRSV ORF). Most preferably, the present protein has a sequence selected from the group consisting of SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:61, SEQ ID NO:67, SEQ ID NO:69 and SEQ ID NO:71.

The present invention may also concern a biologically pure virus, characterized in that it contains the present polynucleic acid and/or that it causes a porcine reproductive and respiratory disease which may include one or more of the following histological lesions: gross and/or microscopic lung lesions (e.g., lung consolidation), Type II pneumocytes, myocarditis, encephalitis, alveolar

exudate formation and syncytia formation. The phrase "biologically pure" refers to a sample of a virus or infectious agent in which all progeny are derived from a single parent. Usually, a "biologically pure" virus sample is achieved by 3 x plaque purification in cell culture.

In particular, the present biologically pure virus or infectious agent is an isolate of the Iowa strain of porcine reproductive and respiratory syndrome virus, samples of which have been deposited under the terms of the Budapest Treaty at the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A., under the accession numbers VR 2385, VR 2386, VR 2428, VR 2429, VR 2430, VR 2431, VR 2474 and VR 2475.

In addition to the characteristics (a)-(e) described above, the Iowa strain of PRRSV may also be characterized by Northern blots of its mRNA. For example, the Iowa strain of PRRSV may contain either 7 or 9 mRNA's, and may also have deletions or variations in their size. In particular, as will be described in the Experiments below, the mRNA's of the Iowa strain of PRRSV may contain up to four deletions, relative to VR 2385/VR 2386.

The present invention further concerns a composition for protecting a pig from viral infection, comprising an amount of the present vaccine effective to raise an immunological response to a virus which causes a porcine reproductive and respiratory disease in a physiologically acceptable carrier.

An effective amount of the present vaccine is one in which a sufficient immunological response to the vaccine is raised to protect a pig exposed to a virus which causes a porcine reproductive and respiratory disease or related illness. Preferably, the pig is protected to an extent in which from on to all of the adverse physiological symptoms or effects (e.g., lung lesions) of the disease to be prevented are found to be significantly reduced.

The composition can be administered in a single dose, or in repeated doses. Dosages may contain, for example, from 1 to 1,000 micrograms of virus-based antigen (vaccine), but should not contain an amount of virus-based antigen sufficient to result in an adverse reaction or physiological symptoms of infection. Methods are known in the art for determining suitable dosages of active antigenic agent.

The composition containing the present vaccine may be administered in conjunction with an adjuvant or with an acceptable carrier which may prolong or sustain the immunological response in the host animal. An adjuvant is a substance that increases the immunological response to the present vaccine when combined therewith. The adjuvant may be administered at the same time and at the same site as the vaccine or at a different time, for example, as a booster. Adjuvants also may advantageously be administered to the animal in a manner or at a site or location different from the manner, site or location in which the vaccine is administered. Adjuvants include aluminum hydroxide, aluminum potassium sulfate, heat-labile or heat-stable enterotoxin isolated from *Escherichia coli*, cholera toxin or the B subunit thereof, diphtheria toxin, tetanus toxin, pertussis toxin, Freund's incomplete adjuvant, Freund's complete adjuvant, and the like. Toxin-based adjuvants, such as diphtheria toxin, tetanus toxin and pertussis toxin, may be inactivated prior to use, for example, by treatment with formaldehyde.

The present invention also concerns a method of protecting a pig from infection against a virus which causes a porcine reproductive and respiratory disease, comprising administering an effective amount of a vaccine which raises an immunological response against such a virus to a pig in need of protection against infection by such a virus. By "protecting a pig from infection" against a

porcine reproductive and respiratory syndrome virus or infectious agent, it is meant that after administration of the present vaccine to a pig, the pig shows reduced (less severe) or no clinical symptoms (such as fever) associated with the corresponding disease, relative to control (infected) pigs. The clinical symptoms may be quantified (e.g., fever, antibody count, and/or lung lesions), semi-quantified (e.g., severity of respiratory distress), or qualified.

The present invention concerns a system for measuring respiratory distress in affected pigs. The present clinical respiratory scoring system evaluates the respiratory distress of affected pigs by the following scale:

0 =no disease; normal breathing

1 =mild dyspnea and polypnea when the pigs are stressed (forced to breathe in larger volumes and/or at an accelerated rate)

2 =mild dyspnea and polypnea when the pigs are at rest

3 =moderate dyspnea and polypnea when the pigs are stressed

4 =moderate dyspnea and polypnea when the pigs are at rest

5 =severe dyspnea and polypnea when the pigs are stressed

6 =severe dyspnea and polypnea when the pigs are at rest

In the present clinical respiratory scoring system, a score of "0" is normal, and indicates that the pig is unaffected by a porcine reproductive and respiratory disease. A score of "3" indicates moderate respiratory disease, and a score of "6" indicates very severe respiratory disease. An amount of the present vaccine or composition may be considered effective if a group of challenged pigs given the vaccine or composition show a lower average clinical respiratory score than a group of

identically challenged pigs not given the vaccine or composition. (A pig is considered "challenged" when exposed to a concentration of an infectious agent sufficient to cause disease in a non-vaccinated animal.)

Preferably, the present vaccine composition is administered directly to a pig not yet exposed to a virus which causes a reproductive or respiratory disease. The present vaccine may be administered orally or parenterally. Examples of parenteral routes of administration include intradermal, intramuscular, intravenous, intraperitoneal, subcutaneous and intranasal routes of administration.

When administered as a solution, the present vaccine may be prepared in the form of an aqueous solution, a syrup, an elixir, or a tincture. Such formulations are known in the art, and are prepared by dissolution of the antigen and other appropriate additives in the appropriate solvent systems. Such solvents include water, saline, ethanol, ethylene glycol, glycerol, A1 fluid, etc. Suitable additives known in the art include certified dyes, flavors, sweeteners, and antimicrobial preservatives, such as thimerosal (sodium ethylmercurithiosalicylate). Such solutions may be stabilized, for example, by addition of partially hydrolyzed gelatin, sorbitol, or cell culture medium, and may be buffered by methods known in the art, using reagents known in the art, such as sodium hydrogen phosphate, sodium dihydrogen phosphate, potassium hydrogen phosphate and/or potassium dihydrogen phosphate.

Liquid formulations may also include suspensions and emulsions. The preparation of suspensions, for example using a colloid mill, and emulsions, for example using a homogenizer, is known in the art.

Parenteral dosage forms, designed for injection into body fluid systems, require proper isotonicity and pH buffering to the corresponding levels of porcine body

fluids. Parenteral formulations must also be sterilized prior to use.

Isotonicity can be adjusted with sodium chloride and other salts as needed. Other solvents, such as ethanol or propylene glycol, can be used to increase solubility of ingredients of the composition and stability of the solution. Further additives which can be used in the present formulation include dextrose, conventional antioxidants and conventional chelating agents, such as ethylenediamine tetraacetic acid (EDTA).

The present invention also concerns a method of producing the present vaccine, comprising the steps of synthesizing or isolating a polynucleic acid of a PRRS virus (preferably the Iowa strain) encoding an antigenic protein or portion thereof (preferably the viral coat protein), infecting a suitable host cell with the polynucleic acid, culturing the host cell, and isolating the antigenic protein or portion thereof from the culture. Alternatively, the polynucleic acid itself can confer immunoprotective activity to a host animal to which it is administered.

Preferably, the vaccine is collected from a culture medium by the steps of (i) precipitating transfected, cultured host cells, (ii) lysing the precipitated cells, and (iii) isolating the vaccine. Particularly preferably, the host cells infected with the virus or infectious agent are cultured in a suitable medium prior to collecting.

Preferably, after culturing infected host cells, the infected host cells are precipitated by adding a solution of a conventional poly(ethylene glycol) (PEG) to the culture medium, in an amount sufficient to precipitate the infected cells. The precipitated infected cells may be further purified by centrifugation. The precipitated cells are then lysed by methods known to those of ordinary skill in the art. Preferably, the cells are lysed by repeated

freezing and thawing (three cycles of freezing and thawing is particularly preferred). Lysing the precipitated cells releases the virus, which may then be collected, preferably by centrifugation. The virus may be isolated and purified by centrifuging in a CsCl gradient, then recovering the appropriate virus-containing band from the CsCl gradient.

Alternatively, the infected cell culture may be frozen and thawed to lyse the cells. The frozen and thawed cell culture material may be used directly as a live vaccine. Preferably, however, the frozen and thawed cell culture material is lyophilized (for storage), then rehydrated for use as a vaccine.

The culture media may contain buffered saline, essential nutrients and suitable sources of carbon and nitrogen recognized in the art, in concentrations sufficient to permit growth of virus-infected cells. Suitable culture media include Dulbecco's minimal essential medium (DMEM), Eagle's minimal essential medium (MEM), Ham's medium, medium 199, fetal bovine serum, fetal calf serum, and other equivalent media which support the growth of virus-infected cells. The culture medium may be supplemented with fetal bovine serum (up to 10%) and/or L-glutamine (up to 2 mM), or other appropriate additives, such as conventional growth supplements and/or antibiotics. A preferred medium is DMEM.

Preferably, the present vaccine is prepared from a virus or infectious agent cultured in an appropriate cell line. The cell line is preferably PSP-36 or an equivalent cell line capable of being infected with the virus and cultured. An example of a cell line equivalent to PSP-36 is the cell line PSP-36-SAH, which was deposited under the terms of the Budapest Treaty at the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A., on October 28, 1992, under the deposit number CRL 11171. Another equivalent cell line is MA-104,

available commercially from Whittaker Bioproducts, Inc. (Walkersville, Maryland). Preliminary results indicate that the Iowa strain of PRRSV can also be cultured in porcine turbinate cells.

There also appears to be a relationship between the severity of histopathology caused by a challenge with a standard amount of a particular isolate and the titer to which the isolate can be grown in a mammalian host cell (e.g., CRL 11171, MA-104 cells [from African green monkey kidney], etc.).

Accordingly, the present invention also concerns a method of culturing a PRRS virus, comprising infecting cell line PSP-36, CRL 11171 or an equivalent cell line and culturing the infected cell line in a suitable medium. An "equivalent cell line" to PSP-36 or CRL 11171 is one which is capable of being infected with the virus and cultured, thereby producing culturable infected cells. Equivalent cell lines include MA-104, PSP-36-SAH and MARC-145 cells (available from the National Veterinary Services Laboratory, Ames, Iowa), for example.

Preferably, the virus cultured is at least one isolate of the Iowa strain of PRRSV. Particularly preferably, the present vaccine is prepared from such a culture of the Iowa strain of PRRSV, cultivated in PSP-36 cells, and plaque-purified at least three times.

The cell line MA-104 is obtained from monkey kidney cells, and is epithelial-like. MA-104 cells form a confluent monolayer in culture flasks containing Dulbecco's minimal essential medium and 10% FBS (fetal bovine serum). When the monolayer is formed, the cells are inoculated with a sample of 10% homogenized tissue, taken from an appropriate tissue (such as lung and/or heart) in an infected pig. Preferably, appropriate antibiotics are present, to permit growth of virus and host cells and to

suppress growth and/or viability of cells other than the host cells (e.g., bacteria or yeast).

Both PSP-36 and MA-104 cells grow some isolates of the PRRS virus to high titers (over 10^7 TCID₅₀/ml). PSP-36 and MA-104 cells will also grow the infectious agent associated with the Iowa strain of PRRSV. MA-104 cells also are able to grow rotaviruses, polioviruses, and other viruses.

CL2621 cells are believed to be of non-porcine origin and are epithelial-like, and are proprietary (Boehringer-Ingelheim). By contrast to PSP-36 and MA-104, some samples of the virus which causes PRRS have been unsuccessfully cultured in CL2621 cells (Bautista et al, American Association of Swine Practitioners Newsletter, 4:32, 1992).

The primary characteristics of CL2621 are that it is of non-swine origin, and is epithelial-like, growing in MEM medium. However, Benfield et al (*J. Vet. Diagn. Invest.*, 1992; 4:127-133) have reported that CL2621 cells were used to propagate PRRS virus, but MA-104 cells were used to control polio virus propagation, thus inferring that CL2621 is not the same as MA-104, and that the same cell may not propagate both viruses.

The Iowa strain of PRRSV generally cannot grow in cell lines other than PSP-36, PSP-36-SAH and MA-104. As described above, however, some viruses which cause PRRS have been reported to grow in both CL2621 and primary swine alveolar macrophages, although some strains of PRRS virus do not grow in PSP-36, MA-104 or CL2621 cells.

The present vaccine, virus isolates, proteins and polynucleic acids can be used to prepare antibodies which may provide immunological resistance to a patient (in this case, a pig) exposed to a virus or infectious agent. Antibodies encompassed by the present invention immunologically bind either to (1) a vaccine which protects a pig against a PRRS virus or (2) to the PRRS virus itself. The present antibodies also have the following utilities:

(1) as a diagnostic agent for determining whether a pig has been exposed to a PRRS virus or infectious agent, and (2) in the preparation of the present vaccine. The present antibody may be used to prepare an immunoaffinity column by known methods, and the immunoaffinity column can be used to isolate the virus or infectious agent, or a protein thereof.

To raise antibodies to such vaccines or viruses, one immunizes an appropriate host animal, such as a mouse, rabbit, or other animals used for such inoculation, with the protein used to prepare the vaccine. The host animal is then immunized (injected) with one of the types of vaccines described above, optionally administering an immune-enhancing agent (adjuvant), such as those described above. The host animal is preferably subsequently immunized from 1 to 5 times at certain intervals of time, preferably every 1 to 4 weeks, most preferably every 2 weeks. The host animals are then sacrificed, and their blood is collected. Sera is then separated by known techniques from the whole blood collected. The sera contains antibodies to the vaccines. Antibodies can also be purified by known methods to provide immunoglobulin G (IgG) antibodies.

The present invention also encompasses monoclonal antibodies to the present vaccines and/or viruses. Monoclonal antibodies may be produced by the method of Kohler et al (Nature, vol. 256 (1975), pages 495-497). Basically, the immune cells from a whole cell preparation of the spleen of the immunized host animal (described above) are fused with myeloma cells by a conventional procedure to produce hybridomas. Hybridomas are cultured, and the resulting culture fluid is screened against the fluid or inoculum carrying the infectious agent (virus or vaccin). Introducing the hybridoma into the peritoneum of th host animal produces a peritoneal growth of th

hybridoma. Collection of the ascites fluid of the host animal provides a sample of the monoclonal antibody to the infectious agent produced by the hybridoma. Also, supernatant from the hybridoma cell culture can be used as a source of the monoclonal antibody, which is isolated by methods known to those of ordinary skill in the art. Preferably, the present antibody is of the IgG or IgM type of immunoglobulin.

The present invention also concerns a method of treating a pig suffering from a reproductive and respiratory disease, comprising administering an effective amount of an antibody which immunologically binds to a virus which causes a porcine reproductive and respiratory disease or to a vaccine which protects a pig against infection by a porcine reproductive and respiratory virus in a physiologically acceptable carrier to a pig in need thereof.

The present method also concerns a method of diagnosing infection of a pig by or exposure of a herd to a porcine reproductive and respiratory syndrome virus and a diagnostic kit for assaying the same, comprising the present antibody (preferably a monoclonal antibody) and a diagnostic agent which indicates a positive immunological reaction with said antibody (preferably comprising peroxidase-conjugated streptavidin, a biotinylated antibody to a PRRSV protein or antigen and a peroxidase). The present kit may further comprise aqueous hydrogen peroxide, a protease which digests the porcine tissue sample, a fluorescent dye (e.g., 3,3'-diaminobenzidine tetrahydrochloride), and a tissue stain (e.g., hematoxylin).

A diagnosis of PRRS relies on compiling information

from the clinical history of the herd being diagnosed, from serology and pathology of infected pigs, and ultimately, on isolation of the PRRS virus (PRRSV) from the infected herd. Thus, the present method of detecting PRRSV is useful in diagnosing infection by and/or exposure to the virus in a herd.

Clinical signs vary widely between farms, and thus, are not the most reliable evidence of a definitive diagnosis, except in the case of a severe acute outbreak in naive herds which experience abortion storms, increased numbers of stillborn pigs, and severe neonatal and nursery pig pneumonia. Presently, the most common clinical presentation is pneumonia and miscellaneous bacterial problems in 3-10 week old pigs. However, many PRRSV-positive herds have no apparent reproductive or respiratory problems.

There are some gross lesions that are very suggestive of PRRSV infection in growing pigs. The most consistent experimentally reproducible gross lesion in 3-10 week-old pigs inoculated with several different PRRSV strains is lymphadenopathy. In particular, iliac and mediastinal lymph nodes are often 3-10 times normal size, tan in color, and sometimes cystic. The lymph nodes are not normally hyperemic, such as the lesion/conditions seen in bacterial septicemia.

Three histologic lesions are consistent with PRRSV infection. Interstitial pneumonia is commonly observed and is characterized by septal infiltration with mononuclear cells, type 2 pneumocyte proliferation, and the presence of necrotic cells in the alveolar spaces. Nonsuppurative perivascular myocarditis and hyperplastic lymph nodes are commonly observed in the subacute stages of disease.

The degree of grossly visible pneumonia is strain dependent. In general, the lungs fail to collapse and have a patchy distribution of 10-80% tan-colored consolidation

with irregular borders. Encephalitis is less often observed. Lesions in the fetus and placenta are rarely observed by light microscopy.

However, the percentage of consolidation in the lungs provides a particularly reliable test for infection by PRRSV (i.e., $\geq 10\%$ consolidation at any time from 3 to 10 days post-infection (DPI) is a positive indication of infection), particularly by a high virulence phenotype (hv) virus ($\geq 40\%$ consolidation at any time from 3 to 10 days DPI is a positive indication of infection by an hv PRRSV isolate).

In contrast to histopathology on lung tissue(s), most laboratories are routinely using either an indirect-fluorescent antibody (IFA) test or immunoperoxidase monolayer assay (IPMA) for serum antibody detection. With both the IFA and IPMA, one must subjectively determine endpoints and thus the tests are not automatable. Serum virus (SVN) neutralization tests have also been developed, and ELISA tests are currently used in some research laboratories. Antibodies detected by the IFA test usually appear with 10 days of exposure but may be relatively short-lived, sometimes disappearing within 3 months.

Antibodies detected by ELISA usually appear within 3 weeks, but their duration is unknown. SVN antibodies usually are not detected until 4-5 weeks post exposure. The SVN test is considered less sensitive in acute disease, but improvements have been made in the SVN test using seronegative porcine serum supplementation. SVN titers reportedly are measurable longer than titers in IFA and IPMA, and thus, may be better suited for detection of positive animals in chronically infected herds.

In IFA, infected cells are fixed with acetone and methanol solutions, and antibodies for the convalescent sera of infected pigs are incubated with the infected cells, preferably for about 30 min. at 37°C. A positive

immunological reaction is one in which the antibody binds to the virus-infected cells, but is not washed out by subsequent washing steps (usually 3 X with PBS buffer). A second antibody (an anti-antibody) labeled with a fluorescent reagent (FITC) is then added and incubated, preferably for another 30 min. A positive immunological reaction results in the second antibody binding to the first, being retained after washing, and resulting in a fluorescent signal, which can be detected and semi-quantified. A negative immunological reaction results in little or no binding of the antibody to the infected cell. Therefore, the second, fluorescently-labeled antibody fails to bind, the fluorescent label is washed out, and little or no fluorescence is detected, compared to an appropriate positive control.

IPA and ELISA kits are similar to the IFA kit, except that the second antibody is labeled with a specific enzyme, instead of a fluorescent reagent. Thus, one adds an appropriate substrate for the enzyme bound to the second antibody which results in the production of a colored product, which is then detected and quantified by colorimetry, for example.

Clinicians use antibody titers to determine the appropriate time for vaccination and/or implementation of management or control strategies. Prior to the present invention, serology tests did not provide antibody titer levels adequate or reliable enough to make animal health care decisions. It may have been appropriate to look for a change from seronegative to seropositive status, or for at least a 4-fold increase in titer, as a positive indication of PRRSV infection/exposure. Looking for an increasing percentage of seropositive pigs in a particular age group over time in a herd can be useful to determine where the virus is maintained and actively spreading. Sows infected

in the early 3rd trimester and aborting near term will likely not show increasing titers, however.

Virus isolation (VI) provides a definitive diagnosis, but has some limitations. Virus is rarely isolated from stillborn or autolyzed aborted fetuses. Sows infected early in the last trimester may have transient viremia and not abort until late term. Dead pigs of any age are not the best samples for VI, because the virus does not survive well at room temperature. Tissues should be removed from the carcass, packaged separately, and refrigerated as soon as possible to obtain a viable virus sample.

The best tissues for virus isolation are tonsil, lung, lymph nodes, and spleen. Serum is also an excellent sample for virus isolation, since (a) viremia is often prolonged in growing pigs, (b) the sample is easy to handle, and (c) the sample can be quickly chilled and processed.

Variation between laboratories in the ability to isolate PRRSV is high because the tests, reagents, cell lines, and media used to detect/screen for PRRSV have not been standardized. The efficacy of isolation varies because not all North American strains will grow on each cell line. Frozen tissue-section IFA tests have been used with limited success.

Serum virus neutralization (SVN) tests have also been developed, and ELISA tests are currently used in some research laboratories. Antibodies detected by ELISA usually appear within 3 weeks, but their duration is unknown. SVN antibodies usually are not detected until 4-5 weeks post-exposure. The SVN test is considered less sensitive in acute disease, but improvements have been made in the SVN test using seronegative porcine serum supplementation. SVN titers reportedly are measurable for a longer period of time than titers in IFA and IPMA. Thus, SVN titers may be better suited for detection of positive animals in chronically infected herds.

Prior to the present invention, however, serology tests did not provide antibody titer levels adequate or reliable enough to make animal health care decisions. Looking for an increasing percentage of seropositive pigs in a particular age group over time in a herd can also be useful to determine where the virus is maintained and actively spreading. Sows infected in the early third trimester and aborting near term will likely not show increasing titers, however. Thus, although it may have been appropriate to look for a change from seronegative to seropositive status or for at least a 4-fold increase in titer as a positive indication of PRRSV infection and/or exposure, a need for a more reliable titer-based assay is felt.

Thus, the present invention also concerns a method for detecting PRRSV antigen in tissues. The present diagnostic method, employing an immunoperoxidase test (IPT) preferably on formalin-fixed tissue, appears to be quite useful to confirm the presence of active infection, and may provide a significant and meaningful increase in the reliability of titer-based assays. A section of lungs, tonsils, mediastinal lymph nodes, and tracheobronchial lymph nodes from 26 pigs experimentally inoculated with ATCC VR 2385 PRRSV was examined (see Experiment V below). The virus was detected in 18/26 lungs, 26/26 tonsils, 15/26 mediastinal lymph nodes, and 14/26 tracheobronchial lymph nodes. The pigs in this study were killed over a 28 day period (post-inoculation). The virus was detected in at least one tissue in every pig necropsied up to 10 days post inoculation.

A complete technique for the present immunoperoxidase technique for PRRSV antigen detection in porcine tissues, based on a streptavidin-biotin assay, is described in Example V hereinunder. Briefly, the present method for detecting PRRSV comprises removing endogenous peroxidase

from an isolated porcine tissue sample with aqueous hydrogen peroxide (preferably, a 0.1-5%, and more preferably, 0.1-1.0% solution), then digesting the tissue with sufficient amount of an appropriate protease to expose viral antigens (for example, Protease XIV, Sigma Chemical Company, St. Louis, MO, and more preferably, a 0.001-0.25% aqueous solution thereof). Thereafter, the method further comprises incubating primary monoclonal antibody ascites fluid (preferably diluted in TRIS/PBS by an amount of from 1:10 to 1:100,000, and more preferably, from 1:100 to 1:10,000) with the protease-treated tissue sections in a humidified chamber for a sufficient length of time and at an appropriate temperature to provide essentially complete immunological binding to occur, if it can in fact occur (e.g., 16 hours at 4°C).

One suitable monoclonal antibody for use in the present diagnostic assay is SDOW-17 (available from Dr. David Benfield, South Dakota State Univ.), which recognizes a conserved epitope of the PRRSV nucleocapsid protein (Nelson et al, "Differentiation of U.S. and European isolates of porcine reproductive and respiratory syndrome virus by monoclonal antibodies," *J. Clin. Micro.*, 31:3184-3189 (1993)).

The present method for detecting PRRSV then further comprises incubating biotinylated goat anti-mouse linking antibody (available from Dako Corporation, Carpinteria, CA) with the tissue, followed by incubating peroxidase-conjugated streptavidin with the biotinylated antibody-treated tissue (Zymed Laboratories, South San Francisco, CA). The method then further comprises incubating the peroxidase-conjugated streptavidin-treated tissue with a chromagen, such as 3,3'-diaminobenzidine tetrahydrochloride (available from Vector Laboratories Inc., Burlingame, CA), and finally, staining the treated tissue with hematoxylin.

Particularly when combined with the further diagnostic techniques of histopathology, virus isolation procedures and serology, the present tissue immunoperoxidase antigen detection technique offers a rapid and reliable diagnosis of PRRSV infection.

Other features of the invention will become apparent in the course of the following descriptions of exemplary embodiments, which are given for illustration of the invention, and are not intended to be limiting thereof.

EXPERIMENT I

MOLECULAR CLONING AND NUCLEOTIDE SEQUENCING OF THE 3'-TERMINAL REGION OF VR 2385 (PLAQUE-PURIFIED ISU-12)

(I) Materials and Methods

(A) Virus Propagation and Purification

A continuous cell line, PSP-36, was used to isolate and propagate ISU-12. The ISU-12 virus was plaque-purified 3 times on PSP-36 cells (plaque-purified ISU-12 virus was deposited under the terms and conditions of the Budapest Treaty at the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A., under Accession No. VR 2385). The PSP-36 cells were then infected with the plaque-purified virus. When more than 70% of the infected cells showed cytopathic changes, the culture was frozen and thawed three times. The culture medium was then clarified by low-speed centrifugation at 5,000 X g for 15 min. at 4°C. The virus was then precipitated with 7% PEG-8000 and 2.3% NaCl at 4°C overnight with stirring, and the precipitate was pelleted by centrifugation. The virus pellets were then resuspended in 2 ml of tris-EDTA buffer, and layered on top of a CsCl gradient (1.1245-1.2858 g/ml). After ultracentrifugation at 28,000 rpm for about 8 hours at 20°C, a clear band with a density of 1.15-1.18 g/ml was observed and harvested.

The infectivity titer of this band was determined by IFA, and the titer was found to be 10^6 TCID₅₀/ml. Typical virus particles were also observed by negative staining electron microscopy (EM).

(B) Isolation of Viral RNA

Total RNA was isolated from the virus-containing band in the CsCl gradient with a commercially available RNA isolation kit (obtained from Stratagene). Poly(A) RNA was then enriched by oligo (dT)-cellulose column chromatography according to the procedure described by the manufacturer of the column (Invitrogen).

(C) Construction of VR 2385 cDNA λ library

A general schematic procedure for the construction of a cDNA λ library is shown in Figure 3. First strand cDNA synthesis from mRNA was conducted by reverse transcription using an oligo (dT) primer having a Xho I restriction site. The nucleotide mixture contained normal dATP, dGTP, dTTP and the analog 5-methyl dCTP, which protects the cDNA from restriction enzymes used in subsequent cloning steps.

Second strand cDNA synthesis was then conducted with RNase H and DNA polymerase I. The cDNA termini were blunted (blunt-ended) with T4 DNA polymerase, ligated to EcoR I adaptors with T4 DNA ligase, and subsequently phosphorylated with T4 polynucleotide kinase. The cDNA was digested with Xho I, and the digested cDNA were size-selected on an agarose gel. Digested cDNA larger than 1 kb in size were selected and purified by a commercially available DNA purification kit (GENECLEAN, available from BIO 101, Inc., La Jolla, California).

The purified cDNA was then ligated into lambda phage vector arms, engineered with Xho I and EcoR I cohesive ends. The ligated vector was packaged into infectious lambda phages with lambda extracts. The SURE strain (available from Stratagene) of *E. coli* cells were used for

transfection, and the lambda library was then amplified and titrated in the XL-1 blue cell strain.

(D) Screening the λ Library by Differential Hybridization

A general schematic procedure for identifying authentic clones of the PRRS virus VR 2385 strain by differential hybridization is shown in Figure 4, and is described hereunder. The λ library was plated on XL-1 blue cells, plaques were lifted onto nylon membranes in duplicates, and denatured with 0.5 N NaOH by conventional methodology. Messenger RNA's from both virus-infected PSP-36 cells and non-infected PSP-36 cells were isolated by oligo (dT)-cellulose column chromatography as described by the manufacturer of the column (Invitrogen).

Complementary DNA probes were synthesized from mRNA's isolated from virus-infected PSP-36 cells and normal PSP-36 cells using random primers in the presence of ^{32}P -dCTP according to the procedure described by the manufacturer (Amersham). Two probes (the first synthesized from virus-infected PSP-36 cells, the other from normal, uninfected PSP-36 cells) were then purified individually by Sephadex G-50 column chromatography. The probes were hybridized with the duplicated nylon membranes, respectively, at 42°C in 50% formamide. Plaques which hybridized with the probe prepared from virus infected cells, but not with the probe prepared from normal cells, were isolated. The phagemids containing viral cDNA inserts were rescued by *in vitro* excision with the help of G408 helper phage. The rescued phagemids were then amplified on XL-1 blue cells. The plasmids containing viral cDNA inserts were isolated by Qiagen column chromatography, and were subsequently sequenced.

(E) Nucleotide Sequencing and Sequence Analysis
Plasmids containing viral cDNA inserts were purified by Qiagen column chromatography, and sequenced by Sanger's

dideoxy method with universal and reverse primers, as well as a variety of internal oligonucleotide primers. Sequences were obtained from at least three separate clones. Additional clones or regions were sequenced when ambiguous sequence data were obtained. The nucleotide sequence data were assembled and analyzed independently using two computer software programs, GENEWORKS (IntelliGenetics, Inc., Mountain View, California) and MACVECTOR (International Biotechnologies, Inc., New Haven, Connecticut).

(F) Oligonucleotide Primers

Oligonucleotides were synthesized as single-stranded DNA using an automated DNA synthesizer (Applied Biosystems) and purified by HPLC. Oligonucleotides PP284 (5'-CGGCCGTGTG GTTCTCGCCA AT-3'; SEQ ID NO:1) and PP285 (5'-CCCCATTTCCTCTAGCGAC TG-3'; SEQ ID NO:2) were synthesized for PCR amplification. A DNA probe was generated with these two primers from the extreme 3' end of the viral genome for Northern blot analysis (see discussion below).

Oligonucleotides PP286 (5'-GCCGCGGAAC CATCAAGCAC-3'; SEQ ID NO:3) and PP287 (5'-CAACTTGACG CTATGTGAGC-3'; SEQ ID NO:4) were synthesized for PCR amplification. A DNA probe generated by these two primers was used to further screen the λ library. Oligonucleotides PP288 (5'-GCGGTCTGGA TTGACGACAG-3'; SEQ ID NO:5), PP289 (5'-GACTGCTAGG GCTTCTGCAC-3'; SEQ ID NO:6), PP386 (5'-GCCATTCAGC TCACATAGCG-3'; SEQ ID NO:7), PP286 and PP287 were used as sequencing primers to obtain internal sequences.

(G) Northern Blot Analysis

A specific DNA fragment from the extreme 3' end of the VR 2385 cDNA clone was amplified by PCR with primers PP284 and PP285. The DNA fragment was excised from an agarose gel with a commercially available DNA purification kit (GENECLEAN, obtained from Bio 101), and labeled with ^{32}P -dCTP by random primer extension (using a kit available

from Amersham). Total RNA was isolated from VR 2385-infected PSP-36 cells at 36 hours post-infection, using a commercially available kit for isolation of total RNA according to the procedure described by the manufacturer (Stratagene). VR 2385 subgenomic mRNA species were denatured with 6 M glyoxal and DMSO, and separated on a 1% agarose gel. (Results from a similar procedure substituting a 1.5% agarose gel are described in Experiment II below and are shown in Figure 5.) The separated subgenomic mRNA's were then transferred onto nylon membranes using a POSIBLOT[®] pressure blotter (Stratagene). Hybridization was carried out in a hybridization oven with roller bottles at 42°C and 50% formamide.

RESULTS

(A) Cloning, Identification and Sequencing of VR 2385 3' Terminal Genome

An oligo (dT)-primed cDNA λ library was constructed from a partially purified virus, obtained from VR 2385-infected PSP-36 cells. Problems were encountered in screening the cDNA λ library with probes based on the Lelystad virus sequence. Three sets of primers were prepared. The first set (PP105 and PP106; SEQ ID NOS:8-9) correspond to positions 14577 to 14596 and 14977 to 14995 of the Lelystad genomic sequence, located in the nucleocapsid gene region. The second set (PP106 and PP107, SEQ ID NOS:9-10) correspond to positions 14977 to 14995 and 14054 to 14072 of the Lelystad genomic sequence, flanking ORF's 6 and 7. The third set (PM541 and PM542; SEQ ID NOS:11-12) correspond to positions 11718 to 11737 and 11394 to 11413 of the Lelystad genomic sequence, located in the ORF-1b region.

PP105: 5'-CTCGTCAAGT ATGCCGGT-3' (SEQ ID NO:8)
PP106: 5'-GCCATTCGCC TGA CTGTCA-3' (SEQ ID NO:9)
PP107: 5'-TTGACGAGGA CTTCCGGCTG-3' (SEQ ID NO:10)
PM541: 5'-GCTCTACCTG CAATTCTGTG-3' (SEQ ID NO:11)

PM542: 5'-GTGTATAGGA CCGGCAACCG-3' (SEQ ID NO:12)

All attempts to generate probes by PCR from the VR 2385 infectious agent using these three sets of primers were unsuccessful. After several attempts using the differential hybridization technique, however, the authentic plaques representing VR 2385-specific cDNA were isolated using probes prepared from VR 2385-infected PSP-36 cells and normal PSP-36 cells. The procedures involved in differential hybridization are described and set forth in Figure 4.

Three positive plaques (λ -4, λ -75 and λ -91) were initially identified. Phagemids containing viral cDNA inserts within the λ phage were rescued by *in vitro* excision with the help of G408 helper phages. The inserts of the positive clones were analyzed by restriction enzyme digestion and terminal sequencing. The specificity of the cDNA clones was further confirmed by hybridization with RNA from PSP-36 cells infected with the Iowa strain of PRRSV, but not with RNA from normal PSP-36 cells. A DNA probe was then generated from the 5'-end of clone λ -75 by PCR with primers PP286 and PP287. Further positive plaques (λ -229, λ -268, λ -275, λ -281, λ -323 and λ -345) were identified using this probe. All λ cDNA clones used to obtain the 3'-terminal nucleotide sequences are presented in Fig. 6. At least three separate clones were sequenced to eliminate any mistakes. In the case of any ambiguous sequence data, additional clones and internal primers (PP288, PP289, PP286, PP287 and PP386) were used to determine the sequence. The 2062-bp 3'-terminal sequence (SEQ ID NO:13) and the amino acid sequences encoded by ORF's 5, 6 and 7 (SEQ ID NOS:15, 17 and 19, respectively) are presented in Figure 7.

(B) A Nested Set of Subgenomic mRNA

Total RNA from virus-infected PSP-36 cells was separated on 1% glyoxal/DMSO agarose gel, and blotted onto nylon membranes. A cDNA probe was generated by PCR with a set of primers (PP284 and PP285) flanking the extreme 3'-terminal region of the viral genome. The probe contains a 3'-nontranslational sequence and most of the ORF-7 sequence. Northern blot hybridization results show that the pattern of mRNA species from PSP-36 cells infected with the Iowa strain of PRRSV is very similar to that of Lelystad virus (LV), equine arteritis virus (EAV), lactate dehydrogenase-elevating virus (LDV) and coronavirus, in that virus replication required the formation of subgenomic mRNA's.

The results also indicate that VR 2385-specific subgenomic mRNA's represent a 3'-nested set of mRNA's, because the Northern blot probe represents only the extreme 3' terminal sequence. The size of VR 2385 viral genomic RNA (14 kb) and 6 subgenomic mRNA's (RNA 2 (3.0 kb), RNA 3 (2.5 kb), RNA 4 (2.2 kb), RNA 5 (1.8 kb), RNA 6 (1.3 kb) and RNA 7 (0.98 kb)) resemble those of LV, although there are differences in both the genome and in subgenomic RNA species. Differences were also observed in the relative amounts of the subgenomic mRNA's, RNA 7 being the most predominant subgenomic mRNA.

(C) Analysis of Open Reading Frames Encoded by Subgenomic RNA

Three large ORF's have been found in SEQ ID NO:13: ORF-5 (nucleotides [nt] 426-1025; SEQ ID NO:14), ORF 6 (nt 1013-1534; SEQ ID NO:16) and ORF 7 (nt 1527-1895; SEQ ID NO:18). ORF 4, located at the 5' end of the resulting sequence, is incomplete in the 2062-bp 3'-terminal sequence of SEQ ID NO:13. ORF'S 5, 6 AND 7 each have a coding capacity of more than 100 amino acids. ORF 5 and ORF 6 overlap each other by 13 bp, and ORF 6 and ORF 7 overlap

each other by 8 bp. Two smaller ORF's located entirely within ORF 7 have also been found, coding for only 37 aa and 43 aa, respectively. Another two short ORF's overlap fully with ORF 5. The coding capacity of these two ORF's is only 29 aa and 44 aa, respectively. No specific subgenomic mRNA's were correlated to these smaller ORF's by Northern blot analysis. ORF 6 and ORF 7 are believed to encode the viral membrane protein and capsid protein, respectively.

(D) Consensus Sequence for Leader Junction

Sequence analysis shows that a short sequence motif, AACC, may serve as the site in the subgenomic mRNA's where the leader is added during transcription (the junction site). The junction site of ORF 6 is found 21 bp upstream from the ATG start codon, and the junction site of ORF 7 is found 13 bp upstream from the ATG start codon, respectively. No AACC consensus sequence has been identified in ORF 5, although it has been found in ORF 5 of LV. Similar junction sequences have been found in LDV and EAV.

(E) 3'-Nontranslational Sequence and Poly (A) Tail

A 151 nucleotide-long (151 nt) nontranslational sequence following the stop codon of ORF 7 has been identified in the genome of VR 2385, compared to 114 nt in LV, 80 nt in LDV and 59 nt in EAV. The length of the poly (A) tail is at least 13 nucleotides. There is a consensus sequence, CCGG/AAATT-poly (A) among PRRS virus VR 2385, LV and LDV in the region adjacent to the poly (A) tail.

(F) Sequence Comparison of VR 2385 and LV Genomes Among ORF's 5, 6 and 7, and Among the Nontranslational Sequences

A comparison of the ORF-5 regions of the genomes of VR 2385 and of the Lelystad virus (SEQ ID NO:20) is shown in Figure 8. The corresponding comparisons of the ORF-6 region, the ORF-7 region, and the nontranslational sequences of VR 2385 (SEQ ID NOS:16, 18 and 22,

respectively) with the corresponding regions of LV (SEQ ID NOS:23, 25 and 27, respectively) are shown in Figures 9, 10 and 11, respectively.

The results of the comparisons are presented in Table 1 below. The nucleotide sequence homologies between LV and VR 2385 of the ORF 5, ORF 6, ORF 7 and the nontranslational sequences are 53%, 78%, 58% and 58%, respectively.

The size of ORF 7 in LV is 15 nt larger than that in VR 2385. Also, the 3'-terminal nontranslational sequence is different in length (150 nt in VR 2385, but only 114 nt in LV). Like LV, the junction sequence, AACC, has also been identified in the genome of the Iowa strain of PRRS virus isolate VR 2385, except for ORF 5. The junction sequence of ORF 6 in VR 2385 is 21 nt upstream from the ATG start codon, whereas the junction sequence of ORF 6 is 28 nt upstream from ATG in LV.

Table 1: Comparison of genes of U.S. PRRSV isolate ATCC VR 2385 with those of European isolate Lelystad virus*

Gene	RNA	Estimated RNA size (in Kb)	ORFs	VR 2385			Lelystad			Homology between VR 2385 & Lelystad
				Size amino acids	N-glycosylation sites	Pred. protein size (kd)	Size amino acids	N-glycosylation sites	Pred. protein size (kd)	
5	5	1.9	5	200	2	22.2	201	2	22.4	53
6	6	1.4	6	174	1	19.1	173	2	18.9	78
7	7	0.9	7	123	2	13.6	128	1	13.8	58
NTR	-	-	-	151 (nt)	-	NA	114 (nt)	0	NA	58 (nt)

*: Based on data presented by Conzelmann et al, *Virology*, 193, 329-339 (1993), Meulenberq et al, *Virology*, 192, 62-72 (1993), and the results presented herein.

EXPERIMENT II

THE EXPRESSION OF VR 2385 GENES IN INSECT CELLS

(A) Production of Recombinant Baculovirus

The ORF-5, ORF-6 and ORF-7 sequences were individually amplified by PCR using primers based on the VR 2385 (ISU-12) genomic nucleotide sequence. ORF-5 was amplified using the following primers:

5'-GGGGATCCGG TATTTGGCAA TGTGTC-3' (SEQ ID NO:28)

3'-GGGAATTCGC CAAGAGCACC TTTTGTGG-5' (SEQ ID NO:29)

ORF-6 was amplified using the following primers:

5'-GGGGATCCAG AGTTTCAGCG G-3' (SEQ ID NO:30)

3'-GGGAATTCTG GCACAGCTGA TTGAC-5' (SEQ ID NO:31)

-71-

ORF-7 was amplified using the following primers:

5'-GGGGATCCTT GTTAAATATG CC-3' (SEQ ID NO:32)

3'-GGGAATTCAC CACGCATTC-5' (SEQ ID NO:33)

The amplified DNA fragments were cloned into baculovirus transfer vector pVL1393 (available from Invitrogen). One μ g of linearized baculovirus AcMNPV DNA (commercially available from Pharmingen, San Diego, California) and 2 μ g of PCR-amplified cloned cDNA-containing vector constructs were mixed with 50 μ l of lipofectin (Gibco), and incubated at 22°C for 15 min. to prepare a transfection mixture.

One hour after seeding HI-FIVE cells, the medium was replaced with fresh Excell 400 insect cell culture medium (available from JR Scientific Co.), and the transfection mixture was added drop by drop. The resulting mixture was incubated at 28°C for six hours. Afterwards, the transfection medium was removed, and fresh Excell 400 insect cell culture medium was added. The resulting mixture was then incubated at 28°C.

Five days after transfection, the culture medium was collected and clarified. Ten-fold dilutions of supernatants were inoculated onto HI-FIVE cells, and incubated for 60 min. at room temperature. After the inoculum was discarded, an overlay of 1.25% of agarose was applied onto the cells. Incubation at 28°C was conducted for four days. Thereafter, clear plaques were selected and picked using a sterile Pasteur pipette. Each plaque was mixed with 1 ml of Grace's insect medium into a 5 ml snap cap tube, and placed in a refrigerator overnight to release the virus from the agarose. Tubes were centrifuged for 30 minutes at 2000 x g to remove agarose, and the supernatants were transferred into new sterile tubes. Plaque purification steps were repeated three times to avoid

possible wild-type virus contamination. Pure recombinant clones were stored at -80°C for further investigation.

(B) Expression of Recombinant Iowa Strain Infectious Agent Proteins

Indirect immunofluorescence assay and radioimmunoprecipitation tests were used to evaluate expression.

Indirect immunofluorescence assay: Hi-five insect cells in a 24-well cell culture cluster plate were infected with wild-type baculovirus or recombinant baculovirus, or were mock-infected. After 72 hours, cells were fixed and stained with appropriate dilutions of swine anti-VR 2385 polyclonal antibodies, followed by fluorescein isothiocyanate-labelled (FITC-labelled) anti-swine IgG. Immunofluorescence was detected in cells infected with the recombinant viruses, but not in mock-infected cells or cells inoculated with wild-type baculovirus. For example, Figure 12 shows HI-FIVE cells infected with the recombinant baculovirus containing the VR 2385 ORF-7 gene (Baculo.PRRSV.7), which exhibit a cytopathic effect. Similar results were obtained with recombinant baculovirus containing ORF-5 (Baculo.PRRSV.5) and ORF-6 (Baculo.PRRSV.6; data not shown). Figures 13 and 14 show HI-FIVE cells infected with a recombinant baculovirus containing the VR 2385 ORF-6 gene and VR 2385 ORF-7 gene, respectively, stained with swine antisera to VR 2385, followed by fluorescein-conjugated anti-swine IgG, in which the insect cells are producing recombinant Iowa strain viral protein. Similar results were obtained with recombinant baculovirus containing ORF-5.

Radioimmunoprecipitation: Radioimmunoprecipitation was carried out with each recombinant virus (Baculo.PRRSV.5, Baculo.PRRSV.6 and Baculo.PRRSV.7) to further determine the antigenicity and authenticity of the recombinant proteins. HI-FIVE insect cells were mock-

infected, or alternatively, infected with each of the recombinant baculoviruses. Two days after infection, methionine-free medium was added. Each mixture was incubated for two hours, and then proteins labeled with ^{35}S -methionine (Amersham) were added, and the mixture was incubated for four additional hours at 28°C. Radiolabeled cell lysates were prepared by three cycles of freezing and thawing, and the cell lysates were incubated with preimmune or immune anti-VR 2385 antisera. The immune complexes were precipitated with Protein A agarose and analyzed on SDS-PAGE after boiling. X-ray film was exposed to the gels at -80°C, and developed. Bands of expected size were detected with ORF-6 (Figure 15) and ORF-7 (Figure 16) products.

EXPERIMENT III

Summary:

The genetic variation and possible evolution of porcine reproductive and respiratory syndrome virus (PRRSV) was determined by cloning and sequencing the putative membrane protein (M, ORF 6) and nucleocapsid (N, ORF 7) genes of six U.S. PRRSV isolates with differing virulence. The deduced amino acid sequences of the putative M and N proteins from each of these isolates were aligned with the corresponding sequences (to the extent known) of one other U.S. isolate, two European isolates, and other members of the proposed arterivirus group, including lactate dehydrogenase-elevating virus (LDV) and equine arteritis virus (EAV).

The putative M and N genes displayed 96-100% amino acid sequence identity among U.S. PRRSV isolates with differing virulence. However, their amino acid sequences varied extensively from those of European PRRSV isolates, and displayed only 57-59% and 78-81% identity, respectively. The U.S. PRRSV isolates were more closely related to LDV than were the European PRRSV isolates. The

N protein of the U.S. isolates and European isolates shared about 50% and 40% amino acid sequence identity with that of LDV, respectively.

The phylogenetic dendrograms constructed on the basis of the putative M and N genes of the proposed arterivirus group were similar and indicated that both U.S. and European PRRSV isolates were related to LDV and were distantly related to EAV. The U.S. and European PRRSV isolates fell into two distinct groups with slightly different genetic distance relative to LDV. The results suggest that U.S. and European PRRSV isolates represent two different genotypes, and that they may have evolved from LDV at different time periods and have existed separately in U.S. and Europe before their association with PRRS was recognized in swine.

ORF 6 encodes the membrane protein (M) of PRRSV, based on the similar characteristics of the ORF 6 of EAV, ORF 2 of LDV, and the M protein of mouse hepatitis virus and infectious bronchitis virus (Meulenberq et al, *Virology*, 192, 62-72 (1993); Conzelmann et al, *Virology*, 193, 329-339 (1993); Mardassi et al, *Abstr. Conf. Res. Workers in Animal Diseases*, Chicago, IL, p. 43 (1993)). The product of ORF 7, the viral nucleocapsid protein (N), is extremely basic and hydrophilic (Meulenberq et al, *Virology*, 192, 62-72 (1993); Conzelmann et al, *Virology*, 193, 329-339 (1993); Murtaugh et al, *Proc. Allen D. Leman Swine Conference*, Minneapolis, MN, pp. 43-45 (1993); Mardassi et al, *Abstr. Conf. Res. Workers in Animal Diseases*, Chicago, IL, p. 43 (1993)).

The amino acid sequences encoded by ORF's 5, 6 and 7 of U.S. isolate VR 2385 and of the European isolate Lelystad virus (LV) have been compared, and the identity (i.e., the percentage of amino acids in sequence which are the same) between the two viruses is only 54%, 78% and 58%, respectively. Thus, striking genetic differences exist

between the U.S. isolate VR 2385 and the European isolate LV (see U.S. application Serial No. 08/131,625, filed October 5, 1993).

However, the U.S. isolate VR 2385 is highly pathogenic compared to European LV. Thus, PRRSV isolates in North America and in Europe appear to be antigenically and genetically heterogeneous, and different genotypes or serotypes of PRRSV may exist.

To further determine the genetic variation among the PRRSV isolates, the putative M and N genes of five additional U.S. PRRSV isolates with differing virulence were cloned and sequenced. Phylogenetic trees based on the putative M and N genes of seven U.S. PRRSV isolates, two European PRRSV isolates and other members of the proposed arterivirus group, including LDV and EAV, have been constructed.

PRRSV isolates (ISU-12 (VR 2385/VR 2386), ISU-22 (VR 2429), ISU-55 (VR 2430), ISU-79, ISU-1894 and ISU-3927 (VR 2431), each of which is disclosed and described in U.S. application Serial No. 08/131,625, filed October 5, 1993) were isolated from pig lungs obtained from different farms in Iowa during PRRS outbreaks, according to the procedure described in U.S. application Serial No. 08/131,625. A continuous cell line, ATCC CRL 11171, was used to isolate and propagate these viruses. All viruses were biologically cloned by three cycles of plaque purification prior to polynucleic acid sequencing.

Pathogenicity studies in caesarean-derived colostrum-deprived (CDCD) pigs, described in U.S. application Serial No. 08/131,625, showed that VR 2385, VR 2429 and ISU-79 were highly pathogenic, whereas VR 2430, ISU-1894 and VR 2431 were not as pathogenic. For example, VR 2385, VR 2429 and ISU-79 produced from 50 to 80% consolidation of the lung tissues in experimentally-infected five-week-old CDCD pigs necropsied at 10 days post inoculation, whereas VR

2430, ISU-1894 and VR 2431 produced only 10 to 25% consolidation of lung tissues in the same experiment.

Experimental Section:

Monolayers of ATCC CRL 11171 cells were infected with each of the PRRSV isolates at the seventh passage at an m.o.i. of 0.1. Total cellular RNA was isolated from infected cells by the guanidine isothiocyanate method (Sambrook et al, "Molecular Cloning: A Laboratory Manual," 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York (1989)). The quality of RNA from each isolate was determined by Northern blot hybridization (data not shown) with a cDNA probe generated from the extreme 3'-end of the VR 2385 genome by the polymerase chain reaction (PCR) with primers PP284 and PP285 (SEQ ID NOS: 1 AND 2), as described in U.S. Application Serial No. 08/131,625. cDNA was synthesized from total cellular RNA with random primers using reverse transcriptase. The synthesized cDNA was amplified by polymerase chain reaction (PCR) as described previously (Meng et al, *J. Vet. Diagn. Invest.*, 5, 254-258 (1993)). Primers for RT-PCR were designed on the basis of a sequence in the genome of VR 2385 which resulted in amplification of the entire protein coding regions of the putative M and N genes (5' primer: 5'-GGGGATCCAGAGTTTCAGCGG-3' (SEQ ID NO:30); 3' primer: 5'-GGGAATTCACCACGCATTC-3' (SEQ ID NO:33)). Unique restriction sites (EcoR I and BamH I) at the termini of the PCR products were introduced by conventional methods. A PCR product with the expected size of about 900 bp was obtained from each of the virus isolates. Southern blot hybridization was then used to confirm the specificity of the amplified products.

The ³²P-labelled cDNA probe from VR 2385 hybridized with the RT-PCR products from each of the above virus isolates. The PCR products of the putative M and N genes

from each of the PRRSV isolates were purified and cloned into vector pSK+ (Meng et al, *J. Vet. Diagn. Invest.* 5, 254-258 (1993)). Plasmids containing the full length putative M and N genes were sequenced with an automated DNA Sequencer (obtained from Applied Biosystems, Inc., Foster City California). Three to four cDNA clones from each virus isolate were sequenced with universal and reverse primers, as well as other virus specific sequencing primers (PP288: 5'-GCGGTCTGGATTGACGAC-3' (SEQ ID NO:5) and PP289: 5'-GACTGCTAGGGCTTCTGC-3' (SEQ ID NO:6), each of which is described in application Serial No. 08/131,625, and DP966: 5'-AATGGGGCTTCTCCGG-3' (SEQ ID NO:34)). The sequences were combined and analyzed by the MACVECTOR (International Biotechnologies, Inc.) and GENEWORKS (IntelliGenetics, Inc.) computer programs.

Analysis of the nucleotide sequences encoding the putative M and N proteins of the five U.S. PRRSV isolates indicated that, like LV (Meulenberg et al, *Virology*, 192, 62-72 (1993)) and VR 2385, the putative M and N genes of each of the five additional U.S. isolates overlapped by 8 base pairs (bp). Figure 17 shows the nucleotide sequence of ORF's 6 and 7 of six U.S. PRRSV isolates and of LV, in which the ISU-12 (VR 2385 and VR 2386) nucleotide sequence (SEQ ID NO:35) is shown first, and in subsequent sequences (SEQ ID NOS:36-41), only those nucleotides which are different are indicated. Start codons are underlined and indicated by (+1>), stop codons are indicated by asterisks (*), are indicated by (-), and the two larger deletions in the putative N gene are further indicated by (^).

Figures 18(A)-(B) show the alignment of amino acid sequences of the putative M (Fig. 18(A)) and N (Fig. 18(B)) genes of the proposed arterivirus group, performed with a GENEWORKS program (IntelliGenetics, Inc.), using the following parameters (default values): cost to open a gap is 5, cost to lengthen a gap is 25, minimum diagonal length

is 4, and maximum diagonal offset is 10. The EAV M gene sequence was omitted because the relatively low sequence identity with PRRSV and LDV requires gaps in the alignments. The VR 2385/VR 2386 sequences (SEQ ID NOS:17 and 19) are shown first, and in subsequent sequences (SEQ ID NOS:43, 45, 47, 49, 51, 24, 53, 55, 57, 59, 61 and 26, respectively), only the differences are indicated. Deletions are indicated by (-), and the two larger deletions in the putative N gene are further indicated by (^).

Numerous substitutions in the nucleotide sequence were distributed randomly throughout the M and N genes in each of the five isolates, as compared to VR 2385. Most of the substitutions are third base silent mutations when converted to amino acid sequences (see Fig. 18). Insertions and deletions are found in the nucleotide sequences of the putative M and N genes when comparing the U.S. isolates to LV, but not found among the U.S. isolates (Fig. 17). For example, there are two larger deletions, 15 and 10 nucleotides each, in the putative N gene of the U.S. isolates as compared to the LV N genome (Fig. 17).

The deduced amino acid sequences of the putative M and N genes from the six Iowa strain PRRSV isolates are aligned with the corresponding N sequence of another U.S. isolate, VR 2332 (Murtaugh et al, *Proc. Allen D. Leman Swine Conference*, Minneapolis, MN, pp. 43-45 (1993)); two European PRRSV isolates, LV (Meulenberg et al, *Virology* 192, 62-72 (1993)) and PRRSV isolate 10 (PRRSV-10) (Conzelmann et al, *Virology*, 193, 329-339 (1993)); two LDV strains, LDV-C (Godney et al, *Virology*, 177, 768-771 (1990)) and LDV-P (Kuo et al, *Virus Res.*, 23, 55-72 (1992)); and EAV (Den Boon et al, *J. Virol.*, 65, 2910-2920 (1991)) (Fig. 18).

The amino acid sequences of the putative N gene are highly conserved among the seven U.S. PRRSV isolates (Fig.

18(B)), and displayed 96-100% amino acid sequence identity (Table 1). However, the putative N proteins of the U.S. PRRSV isolates shared only 57-59% amino acid sequence identity with those of the two European isolates (Table 1), suggesting that the U.S. and the European isolates may represent two different genotypes.

The putative M protein of each of the U.S. isolates was also highly conserved, and displayed higher sequence similarity with the M proteins of the two European isolates (Fig. 18(A)), ranging from 78 to 81% amino acid identity (see Table 2 below). The putative N gene of each of the U.S. PRRSV isolates shared 49-50% amino acid sequence identity with that of the LDV strains, whereas the two European PRRSV isolates shared only 40-41% amino acid identity with that of the LDV strains (Table 2).

Two regions of amino acid sequence deletions, "KKSTAPM" (SEQ ID NO:62) and "ASQG" (SEQ ID NO:63), were found in the putative N proteins of each of the seven U.S. PRRSV isolates, as well as the two LDV strains and EAV, when compared to the two European PRRSV isolates (Fig. 18(B)). These results indicated that the U.S. PRRSV isolates are more closely related to LDV than are the European PRRSV isolates, and that PRRSV may have undergone divergent evolution in the U.S. and in Europe before their association with PRRS was recognized in swine (Murtaugh, *Proc. Allen D. Leman Swine Conference, Minneapolis, MN*, pp. 43-45 (1993)).

The European isolates may have diverged from LDV for a longer time than the U.S. isolates, and hence may have evolved first. However, the amino acid sequence identity of the putative M gene between U.S. PRRSV isolates and LDV strains was similar to that between the European PRRSV isolates and LDV strains (Table 2). The putative M and N genes of the U.S. and European isolates of PRRSV shared

Table 2. Pairwise comparison of the amino acid sequences among the putative nucleocapsid and membrane proteins of the proposed arterivirus group

Virus	VIRUS										EAV
	VR2385	ISU-22	ISU-55	ISU-79	ISU-1894	ISU-3927	VR2332	LV	PRRSV-10	LDV-P	LDV-C
VR2385	***	98	96	98	98	96	96	57	57	49	49
ISU-22	99	***	98	100	100	98	98	57	57	49	49
ISU-55	99	100	***	98	98	97	96	59	59	49	49
ISU-79	98	99	99	***	100	98	98	57	57	49	49
ISU-1894	99	100	100	99	***	98	98	57	57	49	49
ISU-3927	96	97	97	97	97	***	96	59	59	49	49
VR2332	N/A	N/A	N/A	N/A	N/A	N/A	***	57	57	50	49
LV	78	79	79	79	79	81	N/A	***	99	41	40
PRRSV-10	78	79	79	79	79	81	N/A	100	***	41	40
LDV-P	50	51	51	51	51	51	N/A	53	53	***	98
LDV-C	49	50	50	50	50	50	N/A	52	52	96	***
EAV	16	16	16	16	16	15	N/A	17	17	16	17

Note. The values in the table are the percentage identity of amino acid sequences. N/A, not available.

^aNucleocapsid protein comparisons are presented in the upper right half and membrane protein comparisons are presented in the lower left half.

only 15-17% and 22-24% amino acid sequence identity with those of EAV, respectively.

The sequence homology of PRRSV with LDV and EAV suggests that these viruses are closely related and may have evolved from a common ancestor (Plagemann et al, supra; Murtaugh, supra). The high sequence conservation between LDV and PRRSV supported the hypothesis that PRRSV may have evolved from LDV and was rapidly adapted to a new host species (Murtaugh, supra). Asymptomatic LDV infection were found in all strains of mice (Murtaugh, supra; Kuo et al, supra). However, many pig farms are infested with wild rodents (Hooper et al, *J. Vet. Diagn. Invest.*, 6, 13-15 (1994)), so it is possible that PRRSV evolved from LDV-infected mice, and was rapidly adapted to a new host, swine.

The evolutionary relationships of PRRSV with other members of the proposed arterivirus group were determined on the basis of the amino acid sequence of the putative M and N genes. Figure 19 shows a phylogenetic tree of the proposed arterivirus group based on the amino acid sequences of the putative M and N genes of this group. The phylogenetic tree for the N gene is essentially the same as that for the M gene. The length of the horizontal lines connecting one sequence to another is proportional to the estimated genetic distance between sequences, as indicated by the numbers given above each line. The UPGMA (unweighted pair group method with arithmetic mean) trees were constructed with a GENWORKS program (IntelliGenetics, Inc.), which first clusters the two most similar sequences, then the average similarity of these two sequences is clustered with the next most similar sequences or subalignments, and the clustering continued in this manner until all sequences/isolates are located in the tree; both trees are unrooted.

The PRRSV isolates fall into two distinct groups. All the U.S. PRRSV isolates thus far sequenced are closely related and form one group. The two European PRRSV isolates are closely related and form another group. Both the U.S. and European PRRSV isolates are related to LDV strains and are distantly related to EAV (Fig. 19).

The evolution patterns for the putative N and M genes also suggest that PRRSV may be a variant of LDV. For example, the genetic distance of the U.S. PRRSV isolates is slightly closer to LDV than the European PRRSV isolates (Fig. 19), again suggesting that the U.S. and European PRRSV may have evolved from LDV at different time periods and existed separately before their association with PRRS was recognized in swine. European PRRSV may have evolved earlier than U.S. PRRSV. It is also possible that the U.S. and European PRRSV could have evolved separately from different LDV variants which existed separately in the U.S. and Europe.

A striking feature of RNA viruses is their rapid evolution, resulting in extensive sequence variation (Koonin et al, *Critical Rev. Biochem. Mol. Biol.*, 28, 375-430 (1993)). Direct evidence for recombination between different positive-strand RNA viruses has been obtained (Lai, *Microbiol. Rev.*, 56, 61-79 (1992)). Western equine encephalitis virus appears to be an evolutionally recent hybrid between Eastern equine encephalitis virus and another alphavirus closely related to Sindbis virus (Hahn et al, *Proc. Natl. Acad. Sci. USA*, 85, 5997-6001 (1988)). Accordingly, the emergence of PRRSV and its close relatedness to LDV and EAV is not surprising. Although the capsid or nucleocapsid protein has been used for construction of evolutionary trees of many positive-strand RNA viruses, proteins with conserved sequence motifs such as RNA-dependent RNA polymerase, RNA replicase, etc., are

typically more suitable for phylogenetic studies (Koonin et al, supra).

EXPERIMENT IV:

CLONING AND SEQUENCING OF cDNA CORRESPONDING TO ORF'S 2, 3 AND 4 OF PRRSV VR 2385.

The region including ORF's 2, 3, and 4 of the genome of the porcine reproductive and respiratory syndrome virus (PRRSV) isolate VR 2385 was cloned and analyzed. To clone the cDNA of PRRSV VR 2385, ATCC CRL 11171 cells were infected with the virus at a m.o.i. of 0.1, and total cellular RNA was isolated using an RNA Isolation Kit (Stratagene). The mRNA fraction was purified through a Poly(A) Quick column (Stratagene), and the purified mRNA was used to generate a cDNA library. A cDNA oligo dT library was constructed in Uni-ZAP XR λ vector using a ZAP-cDNA synthesis kit (Stratagene), according to the supplier's instructions. Recombinant clones were isolated after screening of the library with an ORF 4 - specific hybridization probe (a 240 b.p. PCR product specific for the 3' end of ORF 4; SEQ ID NO:64). Recombinant pSK + contained PRRSV-specific cDNA was excised *in vivo* from positive λ plaques according to the manufacturer's instructions.

Several recombinant plasmids with nested set of cDNA inserts with sizes ranging from 2.3 to 3.9 kb were sequenced from the 5' ends of the cloned fragments. The nucleotide sequence of SEQ ID NO:65 was determined on at least two independent cDNA clones and was 1800 nucleotides in length (Fig. 21). Computer analysis of the nucleotide and the deduced amino acid sequences was performed using GENEWORKS (IntelliGenetics, Inc.) and MACVECTOR (International Biotechnologies, Inc.) programs.

Three partially overlapping ORF's (ORF 2, ORF 3 and ORF 4) were identified in this region. ORF's 2, 3 and 4 comprised nucleotides 12-779 (SEQ ID NO:66), 635-1396 (SEQ ID NO:68) and 1180-1713 (SEQ ID NO:70), respectively, in the sequenced cDNA fragment.

A comparison of DNA sequences of ORF's 2, 3 and 4 of PRRSV VR 2385 with corresponding ORF's of LV virus (SEQ ID NOS:72, 74 and 76, respectively) is presented in Fig. 22. The level of nucleotide sequence identity (homology) was 65% for ORF 2, 64% for ORF 3 and 66% for ORF 4.

The predicted amino acid sequences encoded by ORF's 2-4 of PRRSV VR 2385 (SEQ ID NOS:67, 69 and 71, respectively) and of LV (SEQ ID NOS:73, 75 and 77, respectively) are shown in Fig 23. A comparison of PRRSV VR 2385 and LV shows a homology level of 58% for the protein encoded by ORF 2, 55% for the protein encoded by ORF 3 and 66% for the protein encoded by ORF 4 (see Fig. 23).

EXPERIMENT V

An immunoperoxidase method of detecting PRRSV

Four 3-week-old colostrum-deprived PRRSV negative animals were inoculated intranasally with $10^{5.8}$ TCID₅₀ of PRRSV U.S. isolate ATCC VR 2386 propagated on ATCC CRL 11171 cells. These pigs were housed on elevated woven-wire decks and fed a commercial milk replacer. Two pigs were necropsied at 4 days post inoculation (DPI) and two at 8 DPI.

At the time of necropsy, the right and left lungs of each pig were separated and inflated via the primary bronchus with 45 ml of one of four fixatives and then immersion fixed for 24 hours. The fixatives used in this experiment included 10% neutral buffered formalin, Bouin's solution, HISTOCHOICE (available from Ambresco, Solon, OH), and a mixture containing 4% formaldehyde and 1% glutaraldehyde (4F:1G). The tissues fixed in Bouin's were rinsed in five 30-minute changes of 70% ethyl alcohol after

4 hours fixation in Bouin's. All the tissues were routinely processed in an automated tissue processor beginning in 70% ethyl alcohol. Tissues were processed to paraffin blocks within 48 hours of the necropsy.

Sections of 3 micron thickness were mounted on poly-L-lysine coated glass slides, deparaffinized with two changes of xylene and rehydrated through graded alcohol baths to distilled water. Endogenous peroxidase was removed by three 10-minute changes of 3% hydrogen peroxide. This was followed by a wash-bottle rinse with 0.05 M TRIS buffer (pH 7.6) followed by a 5-minute TRIS bath. Protease digestion was performed on all tissue sections except those fixed in HISTOCHOICE. Digestion was done in 0.05% protease (Protease XIV, available from Sigma Chem., St. Louis, Mo.) in TRIS buffer for 2 minutes at 37°C. Digestion was followed by a TRIS-buffer wash-bottle rinse and then a 5-minute cold TRIS buffer bath. Blocking for 20 minutes was done with a 5% solution of normal goat serum (available from Sigma Chem., St. Louis, Mo.).

The primary antibody used was the monoclonal antibody SDOW-17 (obtained from Dr. David Benfield, South Dakota State Univ.), diluted 1:1000 in TRIS/PBS (1 part TRIS:9 parts PBS (0.01 M, pH 7.2)). The monoclonal antibody SDOW-17 recognizes a conserved epitope on the PRRSV nucleocapsid protein (Nelson et al, *J. Clin. Microbiol.*, 31:3184-3189). The tissue sections were flooded with primary antibody and incubated at 4°C for 16 hours in a humidified chamber. The primary antibody incubation was then followed by a wash-bottle rinse with TRIS buffer, a 5-minute TRIS buffer bath, and then a 5-minute TRIS buffer bath containing 1% normal goat serum. The sections were flooded with biotinylated goat anti-mouse antisera (obtained from Dako Corporation, Carpinteria, CA) for 30 minutes. The linking antibody incubation was followed by three rinses in TRIS buffer, as was done following primary antibody incubation. The

sections were then treated with peroxidase-conjugated streptavidin, diluted 1:200 in TRIS/PBS, for 40 minutes, followed by a TRIS buffer wash-bottle rinse and a 5-minute TRIS buffer bath. The sections were then incubated with freshly-made 3,3'-diaminobenzidine tetrahydrochloride (DAB, obtained from Vector Laboratories Inc., Burlingame, CA) for 8-10 minutes at room temperature, and then rinsed in a distilled water bath for 5 minutes. Counterstaining was done in hematoxylin (available from Shandon, Inc., Pittsburgh, PA), and the sections were rinsed with Scott's Tap Water (10 g MgSO₄ and 2 g NaHCO₃ in 1 liter ultrapure water), then with distilled water. After dehydration, the sections were covered with mounting media, and then a coverslip was applied.

Two negative controls were included. Substitution of TRIS/PBS buffer in place of the primary antibody was done for one control. The other control was done by substituting uninfected, age-matched, gnotobiotic pig lungs for PRRSV-infected lungs.

Histological changes in infected tissues were characterized by moderate multifocal proliferative interstitial pneumonia with pronounced type 2 pneumocyte hypertrophy and hyperplasia, moderate infiltration of alveolar septa with mononuclear cells, and abundant accumulation of necrotic cell debris and mixed inflammatory cells in the alveolar spaces. No bronchial or bronchiolar epithelial damage was observed. However, there was necrotic cell debris in the smaller airway lumina.

Intense and specific staining in the cytoplasm of infected cells was observed in the formalin- and Bouin's-fixed tissues. Staining was less intense and specific in the 4F:1G-fixed tissues. There was poor staining, poor cellular detail, and moderate background staining in the HISTOCHOICE-fixed tissues. Background staining was negligible with the other fixatives. Cellular detail was

superior in the formalin-fixed tissue sections and adequate in the Bouin's- and 4F:1G-fixed tissues.

The labeled antigen was primarily within the cytoplasm of sloughed cells and macrophages in the alveolar spaces (Fig. 24) and within cellular debris in terminal airway lumina (Fig. 25). When compared to sections from the same block stained with hematoxylin and eosin, it was determined that most of the labeled cells were macrophages, and some were likely sloughed pneumocytes. Lesser intensities of staining were observed in mononuclear cells within the alveolar septa and rarely in hypertrophied type 2 pneumocytes.

Using an immunoperoxidase technique on frozen sections, others were able to detect antigen in epithelial cells of bronchioles and alveolar ducts as well as within cells in the alveolar septa and alveolar spaces (Pol et al, "Pathological, ultrastructural, and immunohistochemical changes caused by Lelystad virus in experimentally induced infections of mystery swine disease (synonym: porcine epidemic abortion and respiratory syndrome (PEARS)), " Vet. Q., 13:137-143). We were unable to detect antigen in bronchiolar epithelium using the present immunoperoxidase method.

The present streptavidin-biotin complex (ABC) technique using a PRRSV monoclonal antibody can be modified as needed to identify PRRSV-infected porcine lungs. Both 10% neutral-buffered formalin and Bouin's solution are acceptable fixatives. Protease digestion enhances the antigen detection without destroying cellular detail. This technique is therefore quite useful for the diagnosis of PRRSV-induced pneumonia of pigs, and for detection of PRRSV in lung tissue samples.

EXPERIMENT VI

An immunohistochemical identification of sites of replication of PRRSV

Summary: Four three-week-old caesarian-derived, colostrum-deprived (CDCD) pigs were inoculated intranasally with an isolate of porcine reproductive and respiratory syndrome virus. All inoculated pigs exhibited moderate respiratory disease. Two pigs were necropsied at 4 days post inoculation (PI) and two at 9 days PI. Moderate consolidation of the lungs and severe enlargement of the lymph nodes were noted at necropsy. Moderate perivascular lymphomacrophagic myocarditis was observed. Marked lymphoid follicular hyperplasia and necrosis was observed in the tonsil, spleen, and lymph nodes.

Porcine reproduction and respiratory syndrome virus antigen was detected by the present streptavidin-biotin immunoperoxidase method primarily within alveolar macrophages in the lung and in endothelial cells and macrophages in the heart. Macrophages and dendritic-like cells in the lymph nodes, spleen, tonsil, and thymus stained intensively positive for PRRSV nucleocapsid protein antigen as well.

Experimental section: Four pigs were snatched from the birth canal of a sow that was positive for PRRSV antibody by indirect immunofluorescent antibody (IFA) examination of serum. The pigs were taken to a different site, housed on elevated woven-wire decks and raised on commercial milk replacer. These pigs were bled at 0, 7, 14, and 21 days of age and found to be negative for PRRSV antibody by the IFA test. No PRRSV was isolated from the serum of the pigs or sow using MARC-145 cells (available from National Veterinary Services Laboratory, Ames, Iowa).